

HIGHER HOPF FORMULAE FOR HOMOLOGY VIA GALOIS THEORY

TOMAS EVERAERT, MARINO GRAN, AND TIM VAN DER LINDEN

ABSTRACT. We use Janelidze's Categorical Galois Theory to extend Brown and Ellis's higher Hopf formulae for homology of groups to arbitrary semi-abelian monadic categories. Given such a category $\mathcal{A}$ and a chosen Birkhoff subcategory $\mathcal{B}$ of $\mathcal{A}$, thus we describe the Barr-Beck derived functors of the reflector of $\mathcal{A}$ onto $\mathcal{B}$ in terms of centralization of higher extensions. In case $\mathcal{A}$ is the category $\mathbf{Gp}$ of all groups and $\mathcal{B}$ is the category $\mathbf{Ab}$ of all abelian groups, this yields a new proof for Brown and Ellis's formulae. We also give explicit formulae in the cases of groups vs. k -nilpotent groups, groups vs. k -solvable groups and precrossed modules vs. crossed modules.

CONTENTS

Introduction	1
1. Semi-abelian and homological categories	3
2. Galois structures	5
3. Higher extensions	7
4. Higher central extensions	12
5. Simplicial extensions	18
6. Homology of extensions with respect to centralization	21
7. The Hopf formula for the second homology object	22
8. The higher Hopf formulae	24
9. Examples	27
Index of notations	34
References	34

INTRODUCTION

Generalizing Hopf's formula [23] for the second integral homology group to higher dimensions is a well-studied problem, that still deserves to be better understood from a categorical perspective. Partial results were originally obtained by Conrad [14], Rodicio [35] and Stöhr [36], and the first complete solution—a formula

2000 *Mathematics Subject Classification.* Primary 18G, 20J, 55N35, 18E10.

Key words and phrases. Semi-abelian category, Hopf formula, homology, Galois theory.

The first author's research is financed by a Ph.D. grant of the Institute of Promotion of Innovation through Science and Technology in Flanders (IWT-Vlaanderen). The third author's research is partly financed by the Eduard Čech Center for Algebra and Geometry, Brno, Czech Republic (project number LC505). Both would like to thank the LMPA for its kind hospitality during their stay in Calais.

describing H_n for all n —is due to Brown and Ellis [11]. Their work was recently extended by Donadze, Inassaridze and Porter in the paper [15]. Whereas Brown and Ellis use topological methods, the latter proof is entirely algebraic, and also considers the case of groups vs. k -nilpotent groups instead of just groups vs. abelian groups.

The aim of our present paper is two-fold: giving a conceptual and elementary proof of the higher Hopf formulae, while at the same time placing them in a very general framework. In our opinion, the simplest approach to a formula for H_n is a proof by induction on n . Now even for groups, such an approach naturally leads to the use of categorical methods: the familiar category of groups must be left for more general ones. On the other hand, consequent reasoning along such lines gives a lot of added generality for free.

How general can we go? Just to give an idea: Brown and Ellis's formulae describe H_n for groups vs. abelian groups; Donadze, Inassaridze and Porter add groups vs. k -nilpotent groups for arbitrary k ; we add groups vs. k -solvable groups, Lie algebras vs. abelian Lie algebras, rings vs. zero rings, precrossed modules vs. crossed modules, etc. Furthermore, even for groups, this yields a proof that's less complicated than the existing ones, and essentially amounts to an application of the Hopf formula for the second homology object and some standard diagram chasing arguments. While based on the same basic idea of using higher dimensional extensions and, in particular, higher presentations of an object, the main difference between our method and previous ones is that we can use an inductive argument, because our formula for H_2 also holds in categories of higher extensions.

That such an approach is possible is due to the existence of the appropriate categorical framework, Janelidze, Márki and Tholen's *semi-abelian categories* [29] and Borceux and Bourn's *homological categories* [3]. These were introduced to capture the fundamental homological properties of the categories of groups, rings, Lie algebras, crossed modules etc. much in the same way as abelian categories do for modules over a ring or sheaves of abelian groups. Our work confirms how well the notion of semi-abelian (or homological) category fulfils this promise.

An important ingredient to understanding the higher Hopf formulae is Janelidze's insight that *centralization of higher extensions* yields the objects that occur in these formulae [26, 27]. For instance, consider a group A presented by a double extension $f: q \longrightarrow p$

$$\begin{array}{ccc} F & \longrightarrow & F/K_2 \\ q \downarrow & & \downarrow p \\ F/K_1 & \longrightarrow & A \end{array}$$

where K_1 and K_2 are normal subgroups of F satisfying $A \cong F/K_1 \cdot K_2$ and such that F , F/K_1 and F/K_2 are free groups. Then the third homology group with coefficients in the group of integers $\mathbb{Z}$ is given by

$$H_3(A, \mathbb{Z}) \cong \frac{[F, F] \cap K_1 \cap K_2}{L_2[f]},$$

where the object $L_2[f] = [K_1 \cap K_2, F] \cdot [K_1, K_2]$, a kind of higher commutator, is defined by the *centralization*

$$\begin{array}{ccc} F/L_2[f] & \cdots\dots\dots \triangleright & F/K_2 \\ \downarrow \nabla & & \downarrow \nabla \\ F/K_1 & \longrightarrow \triangleright & A \end{array}$$

of f —the double extension f universally turned into a central one. Of course, such a *double presentation* f always exists: consider the underlying set/free group comonad $\mathbb{G} = (G, \delta, \epsilon)$ and take the diagram

$$\begin{array}{ccc} G^2 A & \xrightarrow{\epsilon_{GA}} & GA \\ G\epsilon_A \downarrow & & \downarrow \epsilon_A \\ GA & \xrightarrow{\epsilon_A} & A \end{array}$$

for f . In the case of groups, Janelidze realized that also the higher Hopf formulae may be interpreted in these terms [27]. As far as we know, our paper is the first attempt to use this idea for proving the formulae. Next to the concept of semi-abelian categories, part of the fundamental theory is provided by the paper [18], where a proof along the same lines is given of the Hopf formula for the second homology object, and some of the needed homological tools are developed.

A categorical theory of central extensions was developed by Janelidze and Kelly in [28] as an application of Janelidze’s Categorical Galois Theory [24]. This theory is modelled on the situation where $\mathcal{A}$ is a variety of universal algebras and $\mathcal{B}$ a given subvariety of $\mathcal{A}$, and allows one to classify the extensions in $\mathcal{A}$ that are *central* with respect to this chosen subvariety $\mathcal{B}$. The idea of relative centrality—which goes back to the work of the Fröhlich school, see e.g., [19, 33, 20]—has for leading example the variety of groups with its subvariety of abelian groups. Janelidze and Kelly’s theory is general enough to include the case where $\mathcal{A}$ is any semi-abelian category and $\mathcal{B}$ is any given Birkhoff subcategory of $\mathcal{A}$. Modelling the notion of centralization of higher extensions in a categorical way, we are forced to extend this theory, but still within the framework of Categorical Galois Theory. The resulting process of centralization of higher extensions then provides us with the higher commutators that occur in the higher Hopf formulae.

In the last section we explain how these commutators may be calculated explicitly, and we do so in some specific cases: groups vs. abelian groups, groups vs. k -nilpotent groups, groups vs. k -solvable groups, precrossed modules vs. crossed modules. In the important example of precrossed modules the homology objects are described by the same formula (see Theorem 9.6): indeed, the so-called Peiffer commutator plays the same role, in the category of precrossed modules, as the usual commutator of normal subgroups does in the category of groups.

1. SEMI-ABELIAN AND HOMOLOGICAL CATEGORIES

As pointed out above, semi-abelian and homological categories were introduced to capture the homological properties of those categories “sufficiently close” to the category $\mathbf{Gp}$ of all groups. In this section we briefly recall their definition and basic properties.

It is important to note that in general, the difference between a semi-abelian category and an abelian one is quite vast: in an abelian category, every morphism may be factored as a cokernel followed by a kernel; any hom-set $\text{Hom}(B, A)$ carries an abelian group structure; binary products and binary coproducts coincide. None of these properties holds true for the category $\mathbf{Gp}$ of groups: the first one because not every subgroup is a normal subgroup, the second one essentially because the pointwise product of two group homomorphisms need not be a homomorphism, and the third one because a group A with $A \times A \cong A + A$ is always trivial. In view of these differences, it is easily understood why the definition of semi-abelian category might sound unfamiliar at first. Nevertheless, the link with the notion of semi-abelian category is simple and precise: a category $\mathcal{A}$ is abelian if and only if both $\mathcal{A}$ and its dual category $\mathcal{A}^{\text{op}}$ are semi-abelian [29].

Definition 1.1. A category $\mathcal{A}$ is *semi-abelian* when it is pointed, Barr exact and Bourn protomodular and has binary coproducts [29]. $\mathcal{A}$ is *homological* [3] when it is pointed, regular and Bourn protomodular.

Of course some explanation is needed. First of all, in a semi-abelian category, all finite limits and colimits exist. In particular, there is a terminal object 1 and an initial object 0, and it is possible to construct finite products and coproducts, equalizers and coequalizers, pullbacks and pushouts. $\mathcal{A}$ being *pointed* means that $0 \cong 1$, i.e., there is a *zero* object: an object that is both initial and terminal. A map is called *zero* when it factors over 0; given any two objects A and B , there is a unique zero map from B to A . This makes it possible to consider kernels and cokernels: given a morphism $f: B \rightarrow A$ in $\mathcal{A}$, a *kernel* $\text{Ker } f: K[f] \rightarrow B$ of f is a pullback of $0 \rightarrow A$ along f and dually, a *cokernel* $\text{Coker } f: A \rightarrow \text{Cok}[f]$ is a pushout of $B \rightarrow 0$ along f .

$\mathcal{A}$ being *Barr exact* means that it is regular, and such that every internal equivalence relation in $\mathcal{A}$ is a kernel pair [1]. We start by commenting on the regularity, and later come back to the other condition. Recall that a morphism is called a *regular epimorphism* when it is a coequalizer of some pair of arrows. Having finite limits and coequalizers of kernel pairs, $\mathcal{A}$ is *regular* when moreover the regular epimorphisms of $\mathcal{A}$ are pullback-stable. In a regular category, image factorizations exist: any morphism $f: B \rightarrow A$ can be factored as a regular epimorphism $B \rightarrow \text{Im}[f]$ followed by a monomorphism $\text{Im}[f] \rightarrow A$ called the *image* of f ; this factorization is unique up to isomorphism. A morphism f such that $\text{Im } f$ is a kernel is called *proper*.

In this pointed and regular context, $\mathcal{A}$ is *Bourn protomodular* [5] if and only if the (regular) *Short Five Lemma* holds: this means that for every commutative diagram

$$\begin{array}{ccccc} K[f_0] & \xrightarrow{\text{Ker } f_0} & B_0 & \xrightarrow{f_0} & A_0 \\ k \downarrow & & \downarrow b & & \downarrow a \\ K[f] & \xrightarrow{\text{Ker } f} & B & \xrightarrow{f} & A \end{array}$$

such that f and f_0 are regular epimorphisms, k and a being isomorphisms entails that b is an isomorphism. This implies that every regular epimorphism is in fact a cokernel (of its kernel). Accordingly, we can define exact sequences as follows.

A sequence of morphisms $(f_i)_{i \in I}$

$$\cdots \longrightarrow A_{i+1} \xrightarrow{f_{i+1}} A_i \xrightarrow{f_i} A_{i-1} \longrightarrow \cdots$$

is called *exact at A_i* if $\text{Im } f_{i+1} = \text{Ker } f_i$. (In particular, then f_{i+1} is proper.) It is called *exact* when it is exact at A_i , for all $i \in I$. Sequence **A** below

$$0 \longrightarrow K \xrightarrow{k} B \xrightarrow{f} A \longrightarrow 0 \quad (\mathbf{A})$$

is exact if and only if it represents (k, f) as a *short exact sequence*: $k = \text{Ker } f$ and $f = \text{Coker } k$.

Finally, under the above assumptions, $\mathcal{A}$ will be Barr exact if and only if the direct image of a kernel along a regular epimorphism is a kernel: given any kernel k and any regular epimorphism f in $\mathcal{A}$, their composition $f \circ k$ is proper—its image $\text{Im } f \circ k$ is a kernel.

Examples of semi-abelian categories include all abelian categories; any variety of Ω -groups: amongst the operations defining it, there is a group operation and a unique constant (the unit of the group operation), in particular, the categories of groups, (non-unital) rings, (pre)crossed modules, Lie algebras over a field, commutative algebras; compact Hausdorff groups; C^* -algebras; the dual of the category of pointed sets.

For some applications, having a homological category instead of a semi-abelian one will be sufficient. For instance, the category of topological groups and of torsion-free abelian groups are homological but not semi-abelian, and we shall meet another example in Section 3 below. Yet, already in this context the basic homological diagram lemma's hold: the *Snake Lemma*, the 3×3 -*Lemma*, etc. [7]

2. GALOIS STRUCTURES

In this section we recall the basic definition of *categorical Galois structure*, which is crucial for the study of higher central extensions. In particular we give a useful sufficient condition for a Galois structure to be *admissible* in the sense of this theory. We refer the reader to the monograph [4] by Borceux and Janelidze, and in particular to its introduction, for the historical background that led to the development of the theory, as well as for the details of several interesting examples of admissible Galois structures.

Definition 2.1. [25] A *Galois structure* $\Gamma = (\mathcal{A}, \mathcal{B}, \mathcal{E}, \mathcal{Z}, I, H)$ consists of two categories, $\mathcal{A}$ and $\mathcal{B}$, an adjunction

$$\mathcal{A} \xrightleftharpoons[H]{I} \mathcal{B},$$

and classes $\mathcal{E}$ and $\mathcal{Z}$ of morphisms of $\mathcal{A}$ and $\mathcal{B}$ respectively, such that:

- (1) $\mathcal{A}$ has pullbacks along arrows in $\mathcal{E}$;
- (2) $\mathcal{E}$ and $\mathcal{Z}$ contain all isomorphisms, are closed under composition and are pullback-stable;
- (3) $I(\mathcal{E}) \subset \mathcal{Z}$;
- (4) $H(\mathcal{Z}) \subset \mathcal{E}$;
- (5) the counit ϵ is an isomorphism;
- (6) each A -component η_A of the unit η belongs to $\mathcal{E}$.

An element of $\mathcal{E}$ is called an *extension*.

Example 2.2. Let $\mathbf{Gp}$ be the category of groups and $\mathbf{Ab}$ its full reflective subcategory of abelian groups:

$$\mathbf{Gp} \begin{array}{c} \xrightarrow{\text{ab}} \\ \perp \\ \xleftarrow{H} \end{array} \mathbf{Ab}.$$

This adjunction determines a Galois structure $\Gamma = (\mathbf{Gp}, \mathbf{Ab}, \mathcal{E}, \mathcal{Z}, \text{ab}, H)$ where H is the inclusion functor, ab is the abelianization functor, and $\mathcal{E}$ and $\mathcal{Z}$ are the classes of surjective homomorphisms in $\mathbf{Gp}$ and in $\mathbf{Ab}$, respectively. For any group A , the A -component of the unit of the adjunction is given by the canonical quotient $\eta_A: A \rightarrow \text{ab}(A) = A/[A, A]$. In particular every η_A belongs to $\mathcal{E}$.

For an object A of $\mathcal{A}$, let us denote $\mathcal{E}(A)$ the full subcategory of the slice category $\mathcal{A}/A$ determined by the arrows $B \rightarrow A$ in $\mathcal{E}$. If $a: A_0 \rightarrow A$ is an arrow, we will write $a^*: \mathcal{E}(A) \rightarrow \mathcal{E}(A_0)$ for the functor that sends an extension $f: B \rightarrow A$ to its pullback a^*f along a . Since maps in $\mathcal{E}$ are pullback-stable, a^* is well-defined.

For every object A of $\mathcal{A}$, the adjunction (I, H) gives rise to an adjunction

$$\mathcal{E}(A) \begin{array}{c} \xrightarrow{I^A} \\ \perp \\ \xleftarrow{H^A} \end{array} \mathcal{Z}(IA),$$

with I^A defined by $I^A f = If$ on arrows f of $\mathcal{E}(A)$, and H^A defined by $H^A x = (\eta_A)^* Hx$ on arrows x of $\mathcal{Z}(IA)$. The unit and counit of this adjunction will be denoted by η^A and ϵ^A , respectively.

Definition 2.3. A Galois structure Γ is called *admissible* if, for every object A of $\mathcal{A}$, the counit $\epsilon^A: I^A H^A \Rightarrow 1_{\mathcal{Z}(IA)}$ is an isomorphism.

$$\begin{array}{ccc} & \xrightarrow{\pi_2} & \\ A \times_{HIA} HX & \xrightarrow{\eta_A \times_{HIA} HX} & HI(A \times_{HIA} HX) \xrightarrow{\epsilon_x^A} HX \\ \pi_1 = H^A x = \eta_A^* Hx \downarrow & & \downarrow HI\pi_1 \quad \downarrow Hx \\ A & \xrightarrow{\eta_A} & HIA = HIA \end{array} \quad (\text{B})$$

Example 2.4. The Galois structure $\Gamma = (\mathbf{Gp}, \mathbf{Ab}, \mathcal{E}, \mathcal{Z}, \text{ab}, H)$ of Example 2.2 is admissible [24]. More generally, any subvariety of a congruence modular variety determines an admissible Galois structure where $\mathcal{E}$ and $\mathcal{Z}$ are the classes of surjective homomorphisms [28].

Definition 2.5. Let $\mathcal{E}$ be a class of pullback-stable regular epimorphisms in a category $\mathcal{A}$. A full replete $\mathcal{E}$ -reflective subcategory $\mathcal{B}$ of $\mathcal{A}$

$$\mathcal{A} \begin{array}{c} \xrightarrow{I} \\ \perp \\ \xleftarrow{H} \end{array} \mathcal{B}$$

is a *strongly $\mathcal{E}$ -Birkhoff* subcategory of $\mathcal{A}$ if, for every $f: B \longrightarrow A$ in $\mathcal{E}$, all arrows in the diagram

$$\begin{array}{ccccc}
 B & & & & \\
 \searrow \scriptstyle f & & \xrightarrow{\scriptstyle \eta_B} & & \\
 & A \times_{HIA} HIB & \longrightarrow & HIB & \\
 & \downarrow & \lrcorner & \downarrow \scriptstyle HIf & \\
 & A & \xrightarrow{\scriptstyle \eta_A} & HIA &
 \end{array}$$

are in $\mathcal{E}$. In particular, the outer diagram is a *regular pushout* in the sense of Bourn (see [8, 12] and Diagram **D** below).

When $\mathcal{A}$ is a regular category and $\mathcal{E}$ is the class of regular epimorphisms in $\mathcal{A}$, the notion of strongly $\mathcal{E}$ -Birkhoff subcategory of $\mathcal{A}$ is stronger than the classical Birkhoff property. This property requires $\mathcal{B}$ to be closed in $\mathcal{A}$ under subobjects and regular quotients or, equivalently, the outer diagram above to be a pushout of regular epimorphisms. A Birkhoff subcategory (in the classical sense) of a variety of universal algebras is the same thing as a subvariety. The notions of Birkhoff subcategory and of strongly $\mathcal{E}$ -Birkhoff subcategory coincide as soon as $\mathcal{A}$ is exact and satisfies the Mal'tsev property [12]: every internal reflexive relation in $\mathcal{A}$ is an equivalence relation. For instance, if $\mathcal{A}$ is a Mal'tsev variety, then the subvarieties of $\mathcal{A}$ are strongly (regular epi)-Birkhoff subcategories. Recall that every semi-abelian category is exact Mal'tsev [6, 3].

In general, a Galois structure with the property that the adjunction satisfies the strongly $\mathcal{E}$ -Birkhoff property is always admissible:

Proposition 2.6. *Let $\Gamma = (\mathcal{A}, \mathcal{B}, \mathcal{E}, \mathcal{Z}, I, H)$ be a Galois structure, where $\mathcal{B}$ is a strongly $\mathcal{E}$ -Birkhoff subcategory of $\mathcal{A}$, with $\mathcal{E}$ a given class of pullback-stable regular epimorphisms. Then Γ is an admissible Galois structure.*

Proof. This is a consequence of the following two facts. On the one hand, we have that η_f^A is an epimorphism for all $f: B \longrightarrow A$ in $\mathcal{E}(A)$, by the strongly $\mathcal{E}$ -Birkhoff property of $\mathcal{B}$. On the other hand, H^A reflects isomorphisms since, again by the strongly $\mathcal{E}$ -Birkhoff property, η_A is a pullback-stable regular epimorphism: see, for instance, Proposition 1.6 in [30]. To see that these facts indeed imply that ϵ_x^A is an isomorphism, for any $x: X \longrightarrow IA$ in $\mathcal{Z}(IA)$, consider the triangular identity $(H^A \epsilon_x^A) \circ \eta_{H^A x}^A = 1_{H^A x}$. Now, $\eta_{H^A x}^A$ is both a split monomorphism and an epimorphism, hence an iso. This implies that $H^A \epsilon_x^A$ is an iso and, finally, ϵ_x^A is an iso, since H^A reflects isomorphisms. $\square$

3. HIGHER EXTENSIONS

We now restrict our attention to the situation where $\mathcal{A}$ is a homological category. We shall be considering higher-dimensional arrows in $\mathcal{A}$; they are the objects of the following categories.

Definition 3.1. Let 2 be the category generated by a single map $\emptyset \longrightarrow \{1\}$. For any n , write 2^n for the n -fold product $2 \times \cdots \times 2$, and denote the functor category $\text{Hom}(2^n, \mathcal{A})$ as $\text{Arr}^n \mathcal{A}$.

Recall that a *regular epimorphism* is a coequalizer of some pair of arrows. As a first approach to higher-dimensional extensions, we could use higher-dimensional regular epimorphisms:

Definition 3.2. Let $\text{Reg}\mathcal{A}$ be the full subcategory of the category of arrows in $\mathcal{A}$ whose objects are the regular epimorphisms. Denote $\text{Reg}^0\mathcal{A} = \mathcal{A}$ and $\text{Reg}^n\mathcal{A} = \text{Reg}(\text{Reg}^{n-1}\mathcal{A})$ for $n \geq 1$. By an *n-fold regular epimorphism* we mean an object of $\text{Reg}^n\mathcal{A}$, i.e., a regular epimorphism between $(n-1)$ -fold regular epimorphisms.

For brevity, we shall say *n-regular epimorphism* instead of *n-fold regular epimorphism*.

Note that a double (= 2-) regular epimorphism of $\mathcal{A}$ may be considered as a commutative square in $\mathcal{A}$ and, in general, an *n-regular epi* as a particular kind of commutative *n-dimensional diagram*. This is a formal consequence of the fact that the functor $(\cdot) \times 2: \text{Cat} \rightarrow \text{Cat}$ is left adjoint to the functor $\text{Hom}(2, \cdot): \text{Cat} \rightarrow \text{Cat}$: thus an object of $\text{Reg}^2\mathcal{A}$, being a functor $2 \rightarrow \text{Hom}(2, \mathcal{A})$, corresponds to a functor $2 \times 2 \rightarrow \mathcal{A}$, i.e., a commutative square in $\mathcal{A}$; by induction, an object of $\text{Reg}^n\mathcal{A}$ can be seen as a functor $2^n \rightarrow \mathcal{A}$.

It is not difficult to show that an object of $\text{Reg}^2\mathcal{A}$ is a pushout square in $\mathcal{A}$, and in general, an object of $\text{Reg}^n\mathcal{A}$ is an *n-dimensional cube* in $\mathcal{A}$ of which all (two-dimensional) faces are pushouts.

When $\mathcal{A}$ is a semi-abelian category, the category $\text{Arr}^n\mathcal{A}$ is of course semi-abelian, as is any category of $\mathcal{A}$ -valued presheaves. On the other hand, while $\text{Reg}^n\mathcal{A}$ is still homological (see [16]), it is no longer semi-abelian. Indeed, it is well known that for a (non-trivial) abelian category $\mathcal{A}$ the category $\text{Reg}\mathcal{A}$ is *not* abelian; hence it cannot be exact, since it is obviously additive. To see this, consider an object A of $\mathcal{A}$ such that $A \neq 0$ and write τ_A for the unique arrow $A \rightarrow 0$. Then the diagram

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \parallel & & \downarrow \\ A & \longrightarrow & 0 \end{array}$$

represents an epimorphism $1_A \rightarrow \tau_A$ in $\text{Reg}\mathcal{A}$ which is not a normal epimorphism (a cokernel). It follows that $\text{Reg}\mathcal{A}$ is not exact whenever $\mathcal{A}$ is a non-trivial abelian category. It is then easily seen that also for $n \geq 2$, $\text{Reg}^n\mathcal{A}$ need not be semi-abelian.

It is well known that when, in a regular category, a commutative square of regular epimorphisms

$$\begin{array}{ccc} B_0 & \xrightarrow{f_0} \twoheadrightarrow & A_0 \\ \downarrow b & & \downarrow a \\ B & \xrightarrow{f} \twoheadrightarrow & A \end{array} \quad (\text{C})$$

is a pullback, it is a pushout. In a regular category, a commutative square of regular epimorphisms is called a *regular pushout* when the comparison map $r: B_0 \rightarrow P$

to a pullback

$$\begin{array}{ccccc}
 B_0 & & & & \\
 \downarrow b & \searrow f_0 & & & \\
 & P & \dashrightarrow & A_0 & \\
 & \downarrow & & \downarrow a & \\
 & B & \twoheadrightarrow & A & \\
 & & f & &
 \end{array} \quad (\mathbf{D})$$

of f along a is a regular epimorphism. An important aspect of these regular pushouts is made clear by the next result.

Proposition 3.3. [8] *Consider, in a homological category, a commutative diagram of exact sequences, such that f , f_0 , b and a are regular epimorphisms:*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & K[f_0] & \twoheadrightarrow & B_0 & \xrightarrow{f_0} & A_0 \longrightarrow 0 \\
 & & \downarrow k & & \downarrow b & & \downarrow a \\
 0 & \longrightarrow & K[f] & \twoheadrightarrow & B & \xrightarrow{f} & A \longrightarrow 0
 \end{array}$$

The right hand square is a regular pushout if and only if k is regular epi. $\square$

For a regular epimorphism of regular epimorphisms that is a regular pushout in $\mathcal{A}$, this means that its kernel is computed degreewise or, equivalently, that degreewise taking kernels in the diagram above induces a morphism k in $\mathcal{A}$ that is an object of $\mathbf{Reg}\mathcal{A}$. In [12] Carboni, Kelly and Pedicchio show that a regular category $\mathcal{A}$ is Barr exact and Mal'tsev if and only if in $\mathcal{A}$, every pushout of regular epimorphisms is a regular pushout. In particular, a semi-abelian category has this property: a pushout of two regular epimorphisms always exists, and it is a regular pushout. On the other hand, the failure of $\mathbf{Reg}^n\mathcal{A}$ to be exact implies that an $(n+2)$ -regular epimorphism is *not* the same as a regular pushout in $\mathbf{Reg}^n\mathcal{A}$, for $n \geq 1$. This difference gives rise to the notion of n -extension.

Definition 3.4. Let $\mathcal{A}$ be a homological category. A *0-fold extension* in $\mathcal{A}$ is an object of $\mathcal{A}$ and a *1-fold extension* is a regular epimorphism in $\mathcal{A}$. An *n -fold extension* is an object (f_0, f) of $\mathbf{Arr}^n\mathcal{A}$ such that all arrows in the induced diagram $\mathbf{D}$ are $(n-1)$ -fold extensions. The n -fold extensions determine a full subcategory $\mathbf{Ext}^n\mathcal{A}$ of $\mathbf{Arr}^n\mathcal{A}$.

Again, we shall say *n -extension* instead of *n -fold extension*.

As an important technical result, we shall need Proposition 3.9, which generalizes Proposition 3.3 to higher extensions. But first we show that some constructions in $\mathbf{Ext}^n\mathcal{A}$ may be performed in $\mathbf{Arr}^n\mathcal{A}$, which will prove very useful later on.

Proposition 3.5. *Let $\mathcal{E}^n$ be the class of $(n+1)$ -extensions in $\mathcal{A}$. Then:*

- (1) *pullbacks in $\mathbf{Ext}^n\mathcal{A}$ along maps in $\mathcal{E}^n$ exist, are pullbacks in $\mathbf{Arr}^n\mathcal{A}$ and, in particular if $n \geq 1$, are computed degreewise in $\mathbf{Ext}^{n-1}\mathcal{A}$;*
- (2) *maps in $\mathcal{E}^n$ are stable under pulling back in $\mathbf{Ext}^n\mathcal{A}$;*
- (3) *$\mathcal{E}^n$ is closed under composition.*

Proof. If $n = 0$ the statements are true because $\mathcal{A}$ is regular. Now consider $n \geq 1$ and suppose that the statements hold for $k < n$.

To prove the first two statements, consider the following pullback in $\text{Arr}^n \mathcal{A}$ of an n -extension (α_0, α) along an arbitrary morphism (f_0, f) .

$$\begin{array}{ccccc}
 & B'_0 & \xrightarrow{f'_0} & A'_0 & \\
 \beta_0 \swarrow & \downarrow & & \searrow \alpha_0 & \\
 B_0 & \xrightarrow{f_0} & A_0 & & \\
 \downarrow b & \downarrow b' & \downarrow a & \downarrow a' & \\
 & B' & \xrightarrow{f'} & A' & \\
 \beta \swarrow & & & \searrow \alpha & \\
 B & \xrightarrow{f} & A & &
 \end{array}$$

Using that $(n-1)$ -extensions are closed under pulling back, we see that β and β_0 are $(n-1)$ -extensions. We are to show that (β_0, β) is an n -extension; then in particular, it is a pullback in $\text{Ext}^n \mathcal{A}$ of (α_0, α) along (f_0, f) .

Pulling back α along a , β along b yields the commutative diagram

$$\begin{array}{ccc}
 B'_0 & \xrightarrow{f'_0} & A'_0 \\
 (\beta_0, b') \downarrow & & \downarrow (\alpha_0, a') \\
 B_0 \times_B B' & \xrightarrow{f_0 \times_f f'} & A_0 \times_A A'.
 \end{array}$$

By assumption, (α_0, a') is an $(n-1)$ -extension; moreover, this diagram is a pullback, hence, by the induction hypothesis, (β_0, b') is an $(n-1)$ -extension. Now b' , as the composite of (β_0, b') with the second projection of the pullback of b along β , is a composite of $(n-1)$ -extensions, hence, by the induction hypothesis, is an $(n-1)$ -extension itself.

To prove the third statement, considering two composable n -extensions

$$\begin{array}{ccccc}
 C_0 & \xrightarrow{g_0} & B_0 & \xrightarrow{f_0} & A_0 \\
 c \downarrow & & b \downarrow & & a \downarrow \\
 C & \xrightarrow{g} & B & \xrightarrow{f} & A
 \end{array}$$

we immediately see that also $f_0 \circ g_0$ and $f \circ g$ are $(n-1)$ -extensions. Since $(n-1)$ -extensions are stable under pulling back, we moreover get that every arrow in the

diagram

$$\begin{array}{ccccc}
C_0 & \xrightarrow{g_0} & B_0 & & \\
\downarrow r_g & & \parallel & & \\
P_g & \xrightarrow{\bar{g}} & B_0 & \xrightarrow{f_0} & A_0 \\
\downarrow \bar{r}_f & \lrcorner & \downarrow r_f & & \parallel \\
P & \xrightarrow{\quad} & P_f & \xrightarrow{\bar{f}} & A_0 \\
\downarrow \bar{a} & \lrcorner & \downarrow \bar{a} & & \downarrow a \\
C & \xrightarrow{g} & B & \xrightarrow{f} & A
\end{array}$$

is an $(n-1)$ -extension, which proves our claim. $\square$

In particular, the kernel $K[f]$ in $\text{Arr}^n \mathcal{A}$ of an $(n+1)$ -extension $f: B \rightarrow A$ is always an n -extension. We are going to show in Proposition 3.9 that the converse is also true. Suppose a map f in $\text{Ext}^n \mathcal{A}$ has a kernel (in $\text{Arr}^n \mathcal{A}$) that is an n -extension; suppose moreover that this map f is regular epi in $\text{Arr}^n \mathcal{A}$: then this map f is in $\mathcal{E}^n$.

Lemma 3.6. *Suppose that $n \geq 0$. Consider, in $\text{Arr}^n \mathcal{A}$, a commutative diagram of exact sequences*

$$\begin{array}{ccccccc}
0 & \longrightarrow & K[f_0] & \twoheadrightarrow & B_0 & \xrightarrow{f_0} & A \longrightarrow 0 \\
& & \downarrow k & & \downarrow b & & \parallel \\
0 & \longrightarrow & K[f] & \twoheadrightarrow & B & \xrightarrow{f} & A \longrightarrow 0.
\end{array}$$

If f and f_0 are $(n+1)$ -extensions, then $k \in \mathcal{E}^n$ if and only if $b \in \mathcal{E}^n$.

Proof. Whenever, in the above diagram, b , f and f_0 are $(n+1)$ -extensions, the right hand side square is an $(n+2)$ -extension, hence k is an $(n+1)$ -extension by Proposition 3.5. On the other hand, by Proposition 8 in [7], if k is a regular epimorphism then so is b . By induction, it is then easy to verify the following: if B is an n -extension, and both f_0 and k are $(n+1)$ -extensions, then also b is in $\mathcal{E}^n$. $\square$

Lemma 3.7. *Suppose that $n \geq 0$. Consider, in $\text{Arr}^n \mathcal{A}$, a commutative diagram of exact sequences, such that f , f_0 , b and a are $(n+1)$ -extensions:*

$$\begin{array}{ccccccc}
0 & \longrightarrow & K[f_0] & \twoheadrightarrow & B_0 & \xrightarrow{f_0} & A_0 \longrightarrow 0 \\
& & \downarrow k & & \downarrow b & & \downarrow a \\
0 & \longrightarrow & K[f] & \twoheadrightarrow & B & \xrightarrow{f} & A \longrightarrow 0.
\end{array}$$

The right hand square is an $(n+2)$ -extension if and only if k is an $(n+1)$ -extension. In particular, if f is an $(n+2)$ -extension, then its kernel $K[f]$ in $\text{Ext}^{n+1} \mathcal{A}$ exists and is computed degreewise.

Proof. It follows from the previous Lemma. $\square$

Lemma 3.8. *Consider a pair of composable arrows $f: B \longrightarrow A$ and $g: C \longrightarrow B$ in $\text{Arr}^n \mathcal{A}$. If $f \circ g$ is an $(n+1)$ -extension and B an n -extension, then f is an $(n+1)$ -extension. $\square$*

Proposition 3.9. *Suppose that $n \geq 0$. Consider, in $\text{Arr}^n \mathcal{A}$, a short exact sequence*

$$0 \longrightarrow K \xrightarrow{k} B \xrightarrow{f} A \longrightarrow 0 \quad (\text{E})$$

such that B is an n -extension. Then f is an $(n+1)$ -extension if and only if K is an n -extension. $\square$

This result shows that for $n \geq 1$, an n -extension is the same thing as a sequence $\mathbf{A}$ in $\text{Ext}^{n-1} \mathcal{A}$ exact in $\text{Arr}^{n-1} \mathcal{A}$. A kernel in $\text{Arr}^{n-1} \mathcal{A}$ between $(n-1)$ -extensions always has a cokernel, and this cokernel is an n -extension. However, in general, an exact sequence in $\text{Ext}^{n-1} \mathcal{A}$ does not determine an extension! Thus it may also be shown that an n -dimensional arrow in $\mathcal{A}$ is an n -extension exactly when it is an *exact n -presentation* in the sense of [15]. Moreover, this allows one to describe extensions in terms of kernels alone.

Corollary 3.10. *Suppose that $n \geq 0$. Consider, in $\text{Arr}^n \mathcal{A}$, an exact fork*

$$R[f] \xrightarrow[\pi_2]{\pi_1} B \xrightarrow{f} A$$

such that B is an n -extension. Then f is an $(n+1)$ -extension if and only if $R[f]$ is an n -extension.

Proof. This is a consequence of Proposition 3.9 and 3.5: it suffices to note that $\text{Ker } f = \pi_2 \circ \text{Ker } \pi_1$. $\square$

Example 3.11. A split epimorphism in $\text{Ext}^n \mathcal{A}$ is an $(n+1)$ -extension.

Notation 3.12. The forgetful functor $\Upsilon: \text{Ext} \mathcal{A} \longrightarrow \mathcal{A}$ that maps an extension $f: B \longrightarrow A$ to the object B has an obvious right adjoint, namely the functor $\Psi: \mathcal{A} \longrightarrow \text{Ext} \mathcal{A}$ that sends an object B of $\mathcal{A}$ to the extension $B \longrightarrow 0$. Composition yields adjunctions

$$\text{Ext}^n \mathcal{A} \xrightleftharpoons[\Psi^n]{\Upsilon^n} \mathcal{A}.$$

4. HIGHER CENTRAL EXTENSIONS

We shall now establish a sequence of Galois structures Γ_n , such that each determines the next one in the following way: Γ_0 is induced by a Birkhoff subcategory $\mathcal{B}$ of a semi-abelian category $\mathcal{A}$, and Γ_n is a structure on $\text{Ext}^n \mathcal{A}$ with class of extensions $\mathcal{E}^n$ and in which the adjunction is given by centralization of n -extensions with respect to Γ_{n-1} .

4.1. Trivial, central and normal extensions. Let $\Gamma = (\mathcal{A}, \mathcal{B}, \mathcal{E}, \mathcal{Z}, I, H)$ be a Galois structure such that $\mathcal{B}$ is an $\mathcal{E}$ -Birkhoff subcategory of $\mathcal{A}$. From now on, we will omit the inclusion H from our notations (and write $\subseteq$ for the functor H). We shall adopt the terminology of [28], calling *trivial*, *central* and *normal* the following types of extensions:

Definition 4.1. Let $f: B \longrightarrow A$ be an extension. One says that f is

- (1) a *trivial extension* (with respect to Γ), when the next square is a pullback;

$$\begin{array}{ccc} B & \xrightarrow{\eta_B} & IB \\ f \downarrow & & \downarrow If \\ A & \xrightarrow{\eta_A} & IA \end{array}$$

- (2) a *central extension*, when there exists an $a: A_0 \rightarrow A$ in $\mathcal{E}(A)$ such that a^*f is trivial;
 (3) a *normal extension*, when the first projection $\pi_1: R[f] \rightarrow B$ (or, equivalently, the second projection π_2) is a trivial extension.

It is clear that every normal extension is central. Γ being admissible implies that, moreover, every trivial extension is normal. To see this, note that f being trivial implies that $R[f] \cong R[If] \times_{IB} B$; but $R[If]$ is in $\mathcal{B}$, because $\mathcal{B}$ is a full replete reflective subcategory of $\mathcal{A}$; hence the admissibility of Γ entails $R[If] \cong IR[f]$.

Example 4.2. Let us again consider Example 2.2 and its Galois structure $\Gamma = (\mathbf{Gp}, \mathbf{Ab}, \mathcal{E}, \mathcal{Z}, \text{ab}, \subseteq)$. One easily sees that the trivial extensions are exactly the surjective homomorphisms of groups $f: B \rightarrow A$ with the property that the restriction $\hat{f}: [B, B] \rightarrow [A, A]$ of f to the derived subgroups is an isomorphism. It was shown in [24] that an extension $f: B \rightarrow A$ that is central with respect to Γ is the same thing as a central extension in the classical sense: its kernel $K[f]$ is contained in the centre $Z(B)$ of B .

4.2. The functors I_n and J_n . Let $\mathcal{A}$ be a semi-abelian category, $\mathcal{B}$ a Birkhoff subcategory of $\mathcal{A}$ and I the reflector $\mathcal{A} \rightarrow \mathcal{B}$, with unit η . Of course, we may also consider I as a functor $\mathcal{A} \rightarrow \mathcal{A}$. Putting $J(A) = K[\eta_A]$ for each object A of $\mathcal{A}$ yields another functor $J: \mathcal{A} \rightarrow \mathcal{A}$ as well as a short exact sequence of functors

$$0 \longrightarrow J \xrightarrow{\mu} 1_{\mathcal{A}} \xrightarrow{\eta} I \longrightarrow 0.$$

From this sequence, we construct a short exact sequence of functors $\mathbf{Arr} \mathcal{A} \rightarrow \mathbf{Arr} \mathcal{A}$ as follows: let f be an extension $B \rightarrow A$ and (π_1, π_2) its kernel pair. Put $J_1[f] = K[J\pi_1]$ and $J_1f = \Psi J_1[f]$.

$$\begin{array}{ccccc} J_1[f] = K[J\pi_1] & \xrightarrow{\text{Ker } J\pi_1} & JR[f] & \xrightarrow{J\pi_1} & JB \\ \downarrow & & \downarrow \mu_{R[f]} & \xrightarrow{J\pi_2} & \downarrow \mu_B \\ K[\pi_1] & \xrightarrow{\text{Ker } \pi_1} & R[f] & \xrightarrow[\pi_2]{\pi_1} & B \end{array} \quad \begin{array}{c} \text{(i)} \\ \mu_B \end{array}$$

Clearly, this defines a functor $J_1: \mathbf{Arr} \mathcal{A} \rightarrow \mathbf{Arr} \mathcal{A}$. Furthermore, let us define $\mu_f^1 = (\mu_B \circ J\pi_2 \circ \text{Ker } J\pi_1, \alpha_A)$, where α_A is the unique arrow from 0 to A , so that $\eta_f^1 = (\rho_f^1, 1_A) = \text{Coker } \mu_f^1$ yields a short exact sequence

$$0 \longrightarrow J_1 \xrightarrow{\mu^1} 1_{\mathbf{Arr} \mathcal{A}} \xrightarrow{\eta^1} I_1 \longrightarrow 0$$

of functors $\mathbf{Arr} \mathcal{A} \rightarrow \mathbf{Arr} \mathcal{A}$. Indeed, μ_f^1 is a monomorphism because both μ_B and $J\pi_2 \circ \text{Ker } J\pi_1$ are monomorphisms: μ_B by assumption, and $J\pi_2 \circ \text{Ker } J\pi_1$ because it is the normalization of the reflexive, hence effective equivalence relation $(JR[f], J\pi_1, J\pi_2)$. Since μ_B is a monomorphism, the square (i) is a pullback, hence

$\mu_{R[f]} \circ \text{Ker } J\pi_1$ is a normal monomorphism, as an intersection of normal monomorphisms. It follows that $\pi_2 \circ \mu_{R[f]} \circ \text{Ker } J\pi_1$, the regular image of $\mu_{R[f]} \circ \text{Ker } J\pi_1$ along π_2 , is normal in B ; hence so is μ_f^1 .

Since $\text{Arr}\mathcal{A}$ is semi-abelian as soon as $\mathcal{A}$ is, we may repeat this process inductively in order to obtain, for each $n \geq 0$, a short exact sequence

$$0 \longrightarrow J_n \xrightarrow{\mu^n} 1_{\text{Arr}^n \mathcal{A}} \xrightarrow{\eta^n} I_n \longrightarrow 0$$

of functors $\text{Arr}^n \mathcal{A} \longrightarrow \text{Arr}^n \mathcal{A}$. Here we put $J_0 = J$ and $I_0 = I$. As in the case $n = 1$, we write $J_n[f]$ for the domain of $J_n f$ and $I_n[f] = B/J_n[f]$ for the domain of $I_n f$, for any n -extension $f: B \longrightarrow A$. Also, we define $\rho_f^n: B \longrightarrow I_n[f]$ via $\eta_f^n = (\rho_f^n, 1_A)$. Hence, for each n -extension $f: B \longrightarrow A$, we have, in $\text{Arr}^{n-1} \mathcal{A}$, a short exact sequence

$$0 \longrightarrow J_n[f] \xrightarrow{\text{Ker } \rho_f^n} B \xrightarrow{\rho_f^n} I_n[f] \longrightarrow 0.$$

4.3. The Galois structures Γ_n . Given a Birkhoff subcategory $\mathcal{B}$ of a semi-abelian category $\mathcal{A}$, we denote $\Gamma_0 = (\mathcal{A}, \mathcal{B}, \mathcal{E}, \mathcal{Z}, I, \subseteq)$ the associated Galois structure: $\mathcal{E}$ and $\mathcal{Z}$ are the classes of regular epimorphisms in $\mathcal{A}$ and in $\mathcal{B}$, respectively. It is well known that for this structure, the central and normal extensions coincide [28].

Let $\text{CExt}_{\mathcal{B}} \mathcal{A} = \text{CExt}_{\mathcal{B}}^1 \mathcal{A}$ denote the full subcategory of $\text{Ext}^1 \mathcal{A}$ whose objects are the Γ_0 -central extensions. Write $\mathcal{E}^1 = \text{Ext}^2 \mathcal{A}$. We are going to show that I_1 (co)restricts to a reflector $\text{Ext} \mathcal{A} \longrightarrow \text{CExt}_{\mathcal{B}} \mathcal{A}$ and that $\text{CExt}_{\mathcal{B}} \mathcal{A}$ is a strongly $\mathcal{E}^1$ -Birkhoff subcategory of $\text{Ext} \mathcal{A}$. In particular, this gives rise to a Galois structure

$$\Gamma_1 = (\text{Ext} \mathcal{A}, \text{CExt}_{\mathcal{B}} \mathcal{A}, \mathcal{E}^1, \mathcal{Z}^1, I_1, \subseteq_1),$$

where $\mathcal{Z}^1$ consists of all 2-extensions in $\text{CExt}_{\mathcal{B}} \mathcal{A}$ and $\subseteq_1$ denotes the inclusion of $\text{CExt}_{\mathcal{B}} \mathcal{A}$ into $\text{Ext} \mathcal{A}$. Inductively, this may be extended to higher extensions: if

$$\Gamma_{n-1} = (\text{Ext}^{n-1} \mathcal{A}, \text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}, \mathcal{E}^{n-1}, \mathcal{Z}^{n-1}, I_{n-1}, \subseteq_{n-1})$$

is the $(n-1)$ -th Galois structure in the sequence, write

$$\text{CExt}_{\mathcal{B}}^n \mathcal{A} = \text{NExt}_{\text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}}(\text{Ext}^{n-1} \mathcal{A}),$$

where $\text{NExt}_{\text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}}$ is the full subcategory of $\text{Ext}^{n-1} \mathcal{A}$ of normal extensions with respect to Γ_{n-1} . Let Γ_n be the structure

$$(\text{Ext}^n \mathcal{A}, \text{CExt}_{\mathcal{B}}^n \mathcal{A}, \mathcal{E}^n, \mathcal{Z}^n, I_n, \subseteq_n),$$

where $\mathcal{E}^n$ and $\mathcal{Z}^n$ are all $(n+1)$ -extensions in $\text{Ext}^n \mathcal{A}$ and $\text{CExt}_{\mathcal{B}}^n \mathcal{A}$, respectively, and $\subseteq_n$ denotes the inclusion $\text{CExt}_{\mathcal{B}}^n \mathcal{A} \longrightarrow \text{Ext}^n \mathcal{A}$. The notation $\text{CExt}_{\text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}}(\text{Ext}^{n-1} \mathcal{A})$ is explained by Proposition 4.5: for all $n \geq 2$, the normal and central extensions with respect to Γ_{n-1} coincide.

Lemma 4.3. *Let $\mathcal{B}$ be a Birkhoff subcategory of a semi-abelian category $\mathcal{A}$. For each $n \geq 1$, $\text{CExt}_{\mathcal{B}}^n \mathcal{A}$ is a strongly $\mathcal{E}^n$ -Birkhoff subcategory of $\text{Ext}^n \mathcal{A}$. The reflector is given by the restriction of I_n to a functor $\text{Ext}^n \mathcal{A} \longrightarrow \text{CExt}_{\mathcal{B}}^n \mathcal{A}$.*

Proof. If we define $\text{CExt}_{\mathcal{B}}^0 \mathcal{A} = \mathcal{B}$, then the case $n = 0$ is true by assumption. Let us then suppose that the Lemma holds for all $0 \leq k \leq n-1$. We prove it for $k = n$.

First of all, $I_n: \text{Arr}^n \mathcal{A} \longrightarrow \text{Arr}^n \mathcal{A}$ must restrict to a functor $\text{Ext}^n \mathcal{A} \longrightarrow \text{CExt}_{\mathcal{B}}^n \mathcal{A}$. Note that $\text{CExt}_{\mathcal{B}}^n \mathcal{A}$ is well defined by the induction hypothesis. Let $f: B \longrightarrow A$ be

an n -extension. By Lemma 3.8, $I_n f$ is an n -extension as well. We must show that it is normal with respect to Γ_{n-1} .

Let (π_1, π_2) denote the kernel pair of f and (π'_1, π'_2) the kernel pair of $I_n f$, computed in $\text{Ext}^{n-1} \mathcal{A}$, or, equivalently, in $\text{Arr}^{n-1} \mathcal{A}$. Consider the following diagram.

$$\begin{array}{ccccccc}
 0 & \longrightarrow & J_{n-1}R[I_n f] & \rightrightarrows & R[I_n f] & \xrightarrow{\eta_{R[I_n f]}^{n-1}} & I_{n-1}R[I_n f] \longrightarrow 0 \\
 & & \downarrow J_{n-1}\pi'_1 & & \downarrow \pi'_1 & \text{(ii)} & \downarrow I_{n-1}\pi'_1 \\
 0 & \longrightarrow & J_{n-1}I_n[f] & \rightrightarrows & I_n[f] & \xrightarrow{\eta_{I_n[f]}^{n-1}} & I_{n-1}I_n[f] \longrightarrow 0
 \end{array}$$

We must prove that the square (ii) is a pullback in $\text{Ext}^{n-1} \mathcal{A}$. By Proposition 3.5, this is equivalent to proving that it is a pullback in $\text{Arr}^{n-1} \mathcal{A}$, taking into account that $\text{CExt}^{n-1} \mathcal{A}$ is strongly $\mathcal{E}^{n-1}$ -Birkhoff in $\text{Ext}^{n-1} \mathcal{A}$. We are going to show that $J_{n-1}\pi'_1 = J_{n-1}\pi'_2$. Since $J_{n-1}\pi'_1$ and $J_{n-1}\pi'_2$ are jointly monic, this implies that $J_{n-1}\pi'_1$ is a monomorphism hence an iso, so that by Theorem 2.3 in [9], (ii) is a pullback.

Since $\text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}$ is a strongly $\mathcal{E}^{n-1}$ -Birkhoff subcategory of $\text{Ext}^{n-1} \mathcal{A}$, J_{n-1} preserves n -extensions, hence the left hand downward pointing arrow in the diagram

$$\begin{array}{ccccc}
 J_{n-1}R[f] & \xrightarrow[J_{n-1}\pi_2]{J_{n-1}\pi_1} & J_{n-1}B & \xrightarrow{\mu_B^{n-1}} & B \\
 \downarrow & & \downarrow & & \downarrow \rho_f^n \\
 J_{n-1}R[I_n f] & \xrightarrow[J_{n-1}\pi'_2]{J_{n-1}\pi'_1} & J_{n-1}I_n[f] & \xrightarrow{\mu_{I_n[f]}^{n-1}} & I_n[f].
 \end{array}$$

is an n -extension; in particular, it is an epimorphism. Furthermore, $\mu_{I_n[f]}^{n-1}$ is a monomorphism. Hence to show that $J_{n-1}\pi'_1 = J_{n-1}\pi'_2$, it suffices to prove

$$\rho_f^n \circ \mu_B^{n-1} \circ J_{n-1}\pi_1 = \rho_f^n \circ \mu_B^{n-1} \circ J_{n-1}\pi_2.$$

But this follows from the fact that $\rho_f^n \circ \mu_B^{n-1}$ factors over $\text{Coeq}(J_{n-1}\pi_1, J_{n-1}\pi_2)$:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & J_n[f] & \rightrightarrows & J_{n-1}B & \longrightarrow & \frac{J_{n-1}B}{J_n[f]} \longrightarrow 0 \\
 & & \parallel & & \downarrow \mu_B^{n-1} & & \downarrow \text{dotted} \\
 0 & \longrightarrow & J_n[f] & \rightrightarrows & B & \xrightarrow{\rho_f^n} & I_n[f] = \frac{B}{J_n[f]} \longrightarrow 0.
 \end{array}$$

Let us then show that $I_n: \text{Ext}^n \mathcal{A} \rightarrow \text{CExt}_{\mathcal{B}}^n \mathcal{A}$ is a left adjoint. Let $h: f \rightarrow g$ be an arrow in $\text{Ext}^n \mathcal{A}$ with g in $\text{CExt}_{\mathcal{B}}^n \mathcal{A}$. Let (π_1, π_2) be the kernel pair of g . The normality of g implies that $J_{n-1}\pi_1$ is an isomorphism. It follows that $J_n[g] = K[J_{n-1}\pi_1] = 0$, hence $I_n g = g$. Consequently, $I_n h$ gives a factorization $I_n f \rightarrow I_n g = g$:

$$\begin{array}{ccc}
 f & \xrightarrow{\eta_f^n} & I_n f \\
 & \searrow h & \downarrow I_n h \\
 & & g,
 \end{array}$$

which is unique, because η_f^n is an epimorphism.

To see that $\mathbf{CExt}_{\mathcal{B}}^n \mathcal{A}$ is a strongly $\mathcal{E}^n$ -Birkhoff subcategory of $\mathbf{Ext}^n \mathcal{A}$, it remains to be shown that J_n preserves $(n+1)$ -extensions: then the strongly $\mathcal{E}^n$ -Birkhoff property follows from Proposition 3.9. Let $(f_0, f): b \rightarrow a$ be an $(n+1)$ -extension. Denote the kernel pair of $f_0: B_0 \rightarrow A_0$ by (π_1, π_2) and the kernel pair of $f: B \rightarrow A$ by (π'_1, π'_2) . Since $\mathbf{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}$ is a strongly $\mathcal{E}^{n-1}$ -Birkhoff subcategory of $\mathbf{Ext}^{n-1} \mathcal{A}$, J_{n-1} preserves n -extensions. Therefore, both $J_{n-1}R[(f_0, f)]$ and $J_{n-1}b$ are n -extensions. It follows that the right hand square in the diagram

$$\begin{array}{ccccc} J_n[f_0] & \longrightarrow & J_{n-1}R[f_0] & \xrightleftharpoons{J_{n-1}\pi_1} & J_{n-1}B_0 \\ \downarrow & & \downarrow J_{n-1}R[(f_0, f)] & & \downarrow J_{n-1}b \\ J_n[f] & \longrightarrow & J_{n-1}R[f] & \xrightleftharpoons{J_{n-1}\pi'_1} & J_{n-1}B \end{array}$$

is a split epi between n -extensions, therefore it is an $(n+1)$ -extension. We can conclude that the left hand downward pointing arrow is an n -extension, hence $J_n(f_0, f)$ is an $(n+1)$ -extension. $\square$

We are now going to prove that the objects of the category $\mathbf{CExt}_{\mathcal{B}}^n \mathcal{A}$ are indeed the n -fold central extensions with respect to the Galois structure Γ_{n-1} . To prove this, we need the next lemma.

Lemma 4.4. *For any $n \geq 0$, the reflector $I_n: \mathbf{Ext}^n \mathcal{A} \rightarrow \mathbf{CExt}_{\mathcal{B}}^n \mathcal{A}$ preserves pullbacks of split epimorphisms along morphisms in $\mathcal{E}^n$.*

Proof. By the Short Five Lemma, an $(n+2)$ -extension (f_0, f) , considered as a square $\mathbf{C}$ in $\mathbf{Ext}^n \mathcal{A}$, is a pullback if and only if its kernel $K[(f_0, f)]$ is an isomorphism in $\mathbf{Arr}^n \mathcal{A}$. Now let f be a split epimorphism in $\mathbf{Ext}^n \mathcal{A}$, and a an $(n+1)$ -extension, in $\mathbf{Ext}^n \mathcal{A}$ and let $(f_0, f): b \rightarrow a$ denote the extension obtained by pulling back f along a .

Consider the diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & K[(f_0, f)] & \rightrightarrows & b & \xrightarrow{(f_0, f)} & a \longrightarrow 0 \\ & & \downarrow & & \downarrow & \text{(iii)} & \downarrow \\ 0 & \longrightarrow & K[I_n(f_0, f)] & \rightrightarrows & I_n b & \xrightarrow{I_n(f_0, f)} & I_n a \longrightarrow 0 \end{array}$$

in $\mathbf{Arr}^{n+1} \mathcal{A}$. By the strongly $\mathcal{E}^n$ -Birkhoff property of I_n , the arrows $b \rightarrow I_n b$ and $a \rightarrow I_n a$ are $(n+2)$ -extensions. Hence the square (iii) is an $(n+3)$ -extension, since (f_0, f) is a split epimorphism. Since an $\mathcal{E}^{n+1}$ -quotient of an isomorphism in $\mathbf{Ext}^n \mathcal{A}$ is an isomorphism—isomorphisms are stable under pushing out—we get that $K[I_n(f_0, f)]$ is iso, and hence the $(n+2)$ -extension $I_n(f_0, f) = (I_n f_0, I_n f)$, considered as a square in $\mathbf{Arr}^n \mathcal{A}$, is a pullback. $\square$

Proposition 4.5. *For any $n \geq 1$, let Γ_n be the Galois structure constructed above.*

- (1) *The central extensions with respect to Γ_n are the normal extensions. Hence the objects in $\mathbf{CExt}_{\mathcal{B}}^{n+1} \mathcal{A}$ are indeed exactly all extensions that are central with respect to Γ_n .*
- (2) *Every central extension that is a split epimorphism is trivial.*

Proof. Let $f: B \rightarrow A$ be a central extension. Since f^*f is a split epimorphism, (1) will follow from (2) if we show that central extensions are pullback-stable in $\text{Ext}^n \mathcal{A}$. Now this latter property is an easy consequence of the fact that $\text{CExt}_{\mathcal{B}}^{n+1} \mathcal{A}$ is a strongly $\mathcal{E}^{n+1}$ -Birkhoff subcategory of $\text{Ext}^{n+1} \mathcal{A}$, which implies that trivial (and hence also central) extensions are pullback-stable.

Suppose that $f: B \rightarrow A$ is a split epic central extension. Its centrality means that there exists an arrow $a: A_0 \rightarrow A$ in $\mathcal{E}^{n+1}$ for which a^*f is trivial. Using this fact and the previous lemma, one concludes that the exterior rectangle and the left hand square in the diagram

$$\begin{array}{ccccc}
 A_0 \times_A B & \xrightarrow{\pi_2} & B & \xrightarrow{\eta_B^n} & I_n B \\
 \pi_1 \downarrow & \lrcorner & \downarrow f & & \downarrow I_n f \\
 A_0 & \xrightarrow{a} & A & \xrightarrow{\eta_A^n} & I_n A
 \end{array}$$

are pullbacks. Since a is a pullback-stable regular epimorphism, it follows that the right hand square is a pullback, and f is a trivial extension. $\square$

Theorem 4.6. *If $\mathcal{A}$ is a semi-abelian category and $\mathcal{B}$ is a Birkhoff subcategory of $\mathcal{A}$, then for every $n \geq 1$, the n -extensions and central n -extensions give rise to a Galois structure Γ_n . This structure is admissible and strongly $\mathcal{E}^n$ -Birkhoff, and provides the corresponding notion of central $(n+1)$ -extensions.* $\square$

Remark 4.7. Like the categories $\text{Reg}^n \mathcal{A}$, the categories $\text{Ext}^n \mathcal{A}$ need not be exact when $n \geq 1$. However, using Proposition 3.5 and Corollary 3.10 one easily deduces from the exactness of $\text{Arr}^n \mathcal{A}$ that every $(n+1)$ -extension is an effective descent morphism in $\text{Ext}^n \mathcal{A}$. (For more details on descent theory, we refer the reader to [30].) Thus the Fundamental Theorem of Categorical Galois Theory from [25] yields, for any given n -extension $a: A_0 \rightarrow A$, a classification of the n -extensions f with codomain A and a^*f trivial. Suppose $\mathcal{A}$ has enough (regular epi)-projective objects; then any $\text{Ext}^n \mathcal{A}$ has enough $\mathcal{E}^n$ -projectives: for every object A in $\text{Ext}^n \mathcal{A}$, an $(n+1)$ -extension $p: P \rightarrow A$ can be chosen such that P is $\mathcal{E}^n$ -projective. Then, since trivial $(n+1)$ -extensions are pullback-stable in $\text{Ext}^n \mathcal{A}$, for all $(n+1)$ -extensions f and a with codomain A , p^*f is trivial as soon as a^*f is. In this case, the Fundamental Galois Theorem yields a classification of the central $(n+1)$ -extensions with a fixed codomain A .

Example 4.8. Let $I_1 = \text{ab}_1$ be the reflection $\text{Ext}^1 \mathbf{Gp} \rightarrow \text{CExt}_{\mathbf{Ab}}^1 \mathbf{Gp}$ sending an extension $f: B \rightarrow A$ of groups to its centralization $\text{ab}_1 f: B/[K[f], B] \rightarrow A$ (see Example 4.2). This reflection gives rise to the Galois structure

$$\Gamma_1 = (\text{Ext}^1 \mathbf{Gp}, \text{CExt}_{\mathbf{Ab}}^1 \mathbf{Gp}, \mathcal{E}^1, \mathcal{Z}^1, \text{ab}_1, \subseteq_1).$$

It was shown by Janelidze [26] that the double extensions that are central with respect to this Galois structure Γ_1 are precisely those extensions $\mathbf{C}$ with the property that $[K[f_0], K[b]] = \{1\}$ and $[K[f_0] \cap K[b], B_0] = \{1\}$. Recently in [22] a similar characterization was obtained, valid in the context of Mal'tsev varieties.

Notation 4.9. Let $f: B \rightarrow A$ be an n -extension. Recall from the definition of the J_n that $J_n f = \Psi J_n[f]$. It is easily seen that $J_n f$ lies, moreover, in the image of Ψ^n . We write $L_n[f]$ for the object defined via $\Psi^n L_n[f] = J_n f$. It follows that only the “top map” of the unit η_f^n at f is not an iso. We denote it by η_f .

5. SIMPLICIAL EXTENSIONS

We recall the semi-abelian definition of Barr-Beck cotriple homology, and we describe how a suitable comonad on $\mathcal{A}$ produces canonical simplicial resolutions of higher extensions in $\mathcal{A}$.

5.1. Homology in semi-abelian categories. In a pointed category, a *chain complex* $C = (d_n: C_n \rightarrow C_{n-1})_{n \in \mathbb{Z}}$ is a sequence of maps satisfying $d_n \circ d_{n+1} = 0$. As in the abelian case, if it exists, the *n-th homology object* of C is

$$H_n C = \text{Cok}[C_{n+1} \rightarrow K[d_n]],$$

the cokernel of the factorization of d_{n+1} over the kernel of d_n . Now for this definition to be of any interest, one demands that the complex C and the underlying category satisfy some additional properties. Recall that a morphism in a pointed and regular category is called *proper* when its image is a kernel. As soon as the ambient category is, moreover, protomodular, homology of *proper* chain complexes—those with boundary operators of which the image is a kernel—is well-behaved: it characterizes exactness of complexes, and any short exact sequence of proper chain complexes induces a long exact homology sequence [18].

This notion of homology may be extended to simplicial objects: one considers the *normalization functor* $N: \mathcal{SA} \rightarrow \text{Ch}\mathcal{A}$ that maps a simplicial object S in a pointed category with pullbacks $\mathcal{A}$ to its *Moore complex* NS , the chain complex with $N_n S = 0$ for $n < 0$, $N_0 S = S_0$,

$$N_n S = \bigcap_{i=0}^{n-1} K[\partial_i: S_n \rightarrow S_{n-1}]$$

and boundary operators $d_n = \partial_n \circ \bigcap_i \text{Ker } \partial_i: N_n S \rightarrow N_{n-1} S$, for $n \geq 1$. Then $H_n S = H_n NS$. As shown in [18, Theorem 3.6], it is easily seen that when $\mathcal{A}$ is semi-abelian, every simplicial object in $\mathcal{A}$ has a proper Moore complex.

Let $\mathcal{A}$ be an arbitrary category and

$$\mathbb{G} = (G: \mathcal{A} \rightarrow \mathcal{A}, \quad \delta: G \Rightarrow G^2, \quad \epsilon: G \Rightarrow 1_{\mathcal{A}})$$

a comonad on $\mathcal{A}$. Recall the axioms of comonad: $\epsilon_{GA} \circ \delta_A = G\epsilon_A \circ \delta_A = 1_{GA}$ and $\delta_{GA} \circ \delta_A = G\delta_A \circ \delta_A$, for any object A of $\mathcal{A}$. Putting

$$\partial_i = G^i \epsilon_{G^{n-i}A}: G^{n+1}A \rightarrow G^n A, \quad \sigma_i = G^i \delta_{G^{n-i}A}: G^{n+1}A \rightarrow G^{n+2}A,$$

for $0 \leq i \leq n$, gives the sequence $(G^{n+1}A)_{n \in \mathbb{N}}$ the structure of a simplicial object $\mathbb{G}A$ of $\mathcal{A}$. It has an augmentation $\epsilon_A: GA \rightarrow A$; the augmented simplicial object $\epsilon_A: \mathbb{G}A \rightarrow A$ is called the *canonical $\mathbb{G}$ -simplicial resolution of A* . The following naturally generalizes Barr-Beck cotriple homology [2] to the semi-abelian context.

Definition 5.1. [18] Let $\mathcal{A}$ be a category equipped with a comonad $\mathbb{G}$ and $\mathcal{B}$ a semi-abelian category. Let $I: \mathcal{A} \rightarrow \mathcal{B}$ be a functor. For $n \geq 1$, the object

$$H_n(A, I)_{\mathbb{G}} = H_{n-1} N I \mathbb{G} A$$

is the *n-th homology object of A (with coefficients in I) relative to the cotriple $\mathbb{G}$* . This defines a functor $H_n(\cdot, I)_{\mathbb{G}}: \mathcal{A} \rightarrow \mathcal{B}$, for every $n \geq 1$.

Example 5.2. (cf. Example 2.2) When $\mathcal{A}$ is the category of groups, $\text{ab}: \text{Gp} \rightarrow \text{Ab}$ the reflector to its Birkhoff subcategory of abelian groups, and $\mathbb{G}$ the underlying set/free group comonad, it is well known that $H_n(A, \text{ab})_{\mathbb{G}}$ is just the *n-th integral homology group* $H_n(A, \mathbb{Z})$ of A (see page II.6.16 of [34]).

5.2. Comonads derived from a suitable adjunction. From now on we shall assume that $\mathcal{A}$ is semi-abelian. We are going to construct simplicial resolutions for higher extensions: to do so, we consider a comonad $\mathbb{G}$ on $\mathcal{A}$ which is derived from an adjunction

$$\mathcal{A} \begin{array}{c} \xrightarrow{U} \\ \xleftarrow[F]{\tau} \end{array} \mathcal{X}, \quad (\mathbf{F})$$

satisfying conditions (1)–(2) below. In this way, for every n , we shall obtain a comonad $\mathbb{G}_n$ on $\text{Arr}^n \mathcal{A}$ such that every n -extension A in $\mathcal{A}$ yields a simplicial resolution $\mathbb{G}_n A$ in the category $\text{Ext}^n \mathcal{A}$ of n -extensions.

Let then $U: \mathcal{A} \rightarrow \mathcal{X}$ be a functor with a left adjoint $F: \mathcal{X} \rightarrow \mathcal{A}$ and write $\epsilon: F \circ U \Rightarrow 1_{\mathcal{A}}$ and $\zeta: 1_{\mathcal{X}} \Rightarrow U \circ F$ for the counit and unit. This induces a comonad $\mathbb{G}$ as follows: $G = F \circ U$, ϵ is the counit and δ is the natural transformation defined by $\delta_A = F \zeta_{UA}$, for every object A of $\mathcal{A}$. Let us now assume that the following conditions (given by Quillen on page II.5.5 of [34]) are satisfied:

- (1) $\epsilon_A: GA \rightarrow A$ is a regular epimorphism for all objects A of $\mathcal{A}$, and
- (2) FX is (regular epi)-projective for all objects X of $\mathcal{X}$.

Then, in particular, $\epsilon_A: GA \rightarrow A$ is a *projective presentation* of A , for every A in $\mathcal{A}$.

Example 5.3. Let $\mathcal{A}$ be semi-abelian and monadic over $\mathcal{X} = \text{Set}$, and $\mathbb{G}$ the induced comonad on $\mathcal{A}$. Condition (1) then follows from Beck's Theorem: actually, ϵ_A is a coequalizer of $G\epsilon_A$ and ϵ_{GA} . And for every set X , FX is free, hence (regular epi)-projective.

Any variety of algebras is monadic over Set , and thus semi-abelian varieties form an important class of examples. A characterization of such varieties is given by Bourn and Janelidze in their paper [10]. In [21], Gran and Rosický characterize semi-abelian categories, monadic over Set .

Definition 5.4. An object of $\text{Arr}^n \mathcal{A}$ is called *extension-projective* when it is projective with respect to the class $\mathcal{E}^n$ of $(n+1)$ -extensions. An n -extension $f: B \rightarrow A$ is called a *presentation* of A when the object B is extension-projective. An n -extension $f: B \rightarrow A$ is called an *n -fold presentation* (or simply *n -presentation*) when the object B is extension-projective and A is an $(n-1)$ -presentation. (A 0-presentation is an object of $\mathcal{A}$.)

Lemma 5.5. *An object $f: B \rightarrow A$ of $\text{Arr}^n \mathcal{A}$ is extension-projective if and only if both B and A are extension-projective.* $\square$

This lemma implies that an object $f: B \rightarrow A$ of $\text{Arr}^n \mathcal{A}$ is extension-projective if and only if all objects in the corresponding n -cube are projective in $\mathcal{A}$.

Proposition 5.6. *Consider an adjunction $\mathbf{F}$ that satisfies conditions (1)–(2). Then, for every $n \geq 0$, there is an induced adjunction*

$$\text{Arr}^n \mathcal{A} \begin{array}{c} \xrightarrow{U_n} \\ \xleftarrow[F_n]{\tau} \end{array} \text{Arr}^n \mathcal{X} \quad (\mathbf{G})$$

defined degreewise, by putting, for objects $f: B \longrightarrow A$ of $\text{Arr}^n \mathcal{A}$ and $z: Y \longrightarrow X$ of $\text{Arr}^n \mathcal{X}$,

$$\begin{aligned} U_n(f: B \longrightarrow A) &= U_{n-1}f: U_{n-1}B \longrightarrow U_{n-1}A; \\ F_n(z: Y \longrightarrow X) &= F_{n-1}z: F_{n-1}Y \longrightarrow F_{n-1}X; \\ \epsilon_n(f: B \longrightarrow A) &= (\epsilon_{n-1}B, \epsilon_{n-1}A): F_n U_n f \longrightarrow f; \\ \zeta_n(z: Y \longrightarrow X) &= (\zeta_{n-1}Y, \zeta_{n-1}X): z \longrightarrow U_n F_n z. \end{aligned}$$

We will often omit the indices and simply write U , F , ϵ and ζ . Putting $G_n = F_n \circ U_n$ and $(\delta_n)_A = F(\zeta_n)_{U_n A}$ for every object A of $\text{Arr}^n \mathcal{A}$ defines a comonad $\mathbb{G}_n = (G_n, \delta_n, \epsilon_n)$, denoted by (G, δ, ϵ) when confusion is unlikely. Moreover, the following two conditions are satisfied:

- (1) $\epsilon_A: GA \longrightarrow A$ is an $(n+1)$ -extension for all n -extensions A , and
- (2) FX is extension-projective for all objects X of $\text{Arr}^n \mathcal{X}$.

Proof. We use induction on n .

Since an object $f: B \longrightarrow A$ of $\text{Arr}^n \mathcal{A}$ is extension-projective if and only if both A and B are extension-projective, condition (2) follows immediately from the induction hypothesis.

In order to show that condition (1) holds as well, it suffices to prove that, for any n -extension $f: B \longrightarrow A$, also $Gf: GB \longrightarrow GA$ is an n -extension. Indeed, if this is the case, then the kernel pair $R[Gf]$ of the arrow Gf is an $(n-1)$ -extension, by Corollary 3.10. Also, $R[f]$ is an $(n-1)$ -extension, being the kernel pair of the n -extension f . Hence, by the induction hypothesis, $\epsilon_{R[f]}$ is an n -extension. Furthermore, by the universal property of the kernel pair $R[Gf]$, there exists an arrow $a: GR[f] \longrightarrow R[Gf]$ such that $\bar{\epsilon}_B \circ a = \epsilon_{R[f]}$, where $\bar{\epsilon}_B$ is the restriction of ϵ_B to $R[Gf] \longrightarrow R[f]$. By Lemma 3.8 this implies that $\bar{\epsilon}_B$ is an n -extension. Finally, by Corollary 3.10, this implies that the right hand square in the next diagram is an $(n+1)$ -extension.

$$\begin{array}{ccccc} GR[f] & & & & \\ \downarrow a & \searrow & & & \\ R[Gf] & \rightrightarrows & GB & \xrightarrow{Gf} & GA \\ \downarrow \bar{\epsilon}_B & & \downarrow \epsilon_B & & \downarrow \epsilon_A \\ R[f] & \rightrightarrows & B & \xrightarrow{f} & A \end{array}$$

Let us then prove that $Gf: GB \longrightarrow GA$ is indeed an n -extension, for any n -extension $f: B \longrightarrow A$.

Let X be any object of $\text{Arr}^{n-1} \mathcal{X}$. Since, by assumption, FX is extension-projective, the map of Hom-sets

$$\text{Hom}(FX, f): \text{Hom}(FX, B) \longrightarrow \text{Hom}(FX, A)$$

is surjective. Since F is left adjoint to U , this implies that the map

$$\text{Hom}(X, Uf): \text{Hom}(X, UB) \longrightarrow \text{Hom}(X, UA)$$

is surjective as well. Since this is the case for any object X of $\text{Arr}^{n-1} \mathcal{X}$, it follows that $Uf: UB \longrightarrow UA$ is a split epimorphism. Consequently, also $Gf: GB \longrightarrow GA$

is a split epimorphism. Since both its domain and codomain are $(n-1)$ -extensions, it is an n -extension (see Example 3.11). $\square$

In particular, $\text{Ext}^n \mathcal{A}$ has enough $\mathcal{E}^n$ -projectives, because for any n -extension A , the map $\epsilon_A: GA \rightarrow A$ is an $(n+1)$ -extension with an $\mathcal{E}^n$ -projective domain.

Corollary 5.7. *If $\mathbb{G}$ is a comonad on a semi-abelian category $\mathcal{A}$ induced by an adjunction $\mathbf{F}$ satisfying conditions (1)–(2), then every object A of $\mathcal{A}$ has a canonical n -presentation: $\epsilon_A^n = \epsilon_{\epsilon_A^{n-1}}: G\epsilon_A^{n-1} \rightarrow \epsilon_A^{n-1}$.* $\square$

6. HOMOLOGY OF EXTENSIONS WITH RESPECT TO CENTRALIZATION

From now on, we concentrate on the situation where

- (1) $\mathcal{A}$ is a semi-abelian category and $\mathcal{B}$ is a Birkhoff subcategory of $\mathcal{A}$;
- (2) $\mathbb{G}$ is a comonad on $\mathcal{A}$ like in Subsection 5.2: $\mathbb{G}$ is derived from an adjunction $\mathbf{F}$ which satisfies the conditions (1)–(2) from Subsection 5.2.

We showed that for all n , there is a Galois structure Γ_n on $\text{Ext}^n \mathcal{A}$ and an $\mathcal{E}^n$ -comonad $\mathbb{G}_n$ on $\text{Arr}^n \mathcal{A}$. Let $f: B \rightarrow A$ be a presentation of an $(n-1)$ -extension A (f may for instance be an n -presentation). In this section, we describe the homology with respect to $\mathcal{B}$ -centralization of f in terms of the homology with respect to $\text{CExt}_{\mathcal{B}}^{n-1} \mathcal{A}$ of A . If I denotes the reflector of $\mathcal{A}$ onto $\mathcal{B}$, and $I_1: \text{Ext} \mathcal{A} \rightarrow \text{CExt}_{\mathcal{B}} \mathcal{A}$ the centralization functor, then for all $k \geq 2$,

$$H_k(f, I_1)_{\mathbb{G}_1} \cong (H_{k+1}(A, I)_{\mathbb{G}} \rightarrow 0),$$

where $\mathbb{G}_1$ is the comonad on $\text{Arr} \mathcal{A}$, induced by $\mathbb{G}$. Observe that the isomorphism above lives in the category $\text{Arr} \mathcal{A}$.

More generally, if I_n denotes the reflector of $\text{Ext}^n \mathcal{A}$ onto $\text{CExt}_{\mathcal{B}}^n \mathcal{A}$, then for all $n \geq 1$ and $k \geq 2$,

$$H_k(f, I_n)_{\mathbb{G}_n} \cong (H_{k+1}(A, I_{n-1})_{\mathbb{G}_{n-1}} \rightarrow 0),$$

where $\mathbb{G}_n$ is the comonad on $\text{Arr}^n \mathcal{A}$, induced by $\mathbb{G}$.

Remark 6.1. For any n , we shall consider I_n as a functor $\text{Ext}^n \mathcal{A} \rightarrow \text{Arr}^n \mathcal{A}$; then the situation fits into the scope of Definition 5.1.

Lemma 6.2. *For every $n \geq 1$, given an n -extension f , there exists an isomorphism $I_{n-1}I_n f \cong I_{n-1}f$ in $\text{Arr}^n \mathcal{A}$.*

Proof. Suppose $f: B \rightarrow A$ is an n -extension. Consider the 3×3 -diagram below. Note that all rows are short exact sequences, as is the middle column.

$$\begin{array}{ccccc} J_n[f] & \xlongequal{\quad} & J_n[f] & \twoheadrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ J_{n-1}B & \twoheadrightarrow & B & \twoheadrightarrow & I_{n-1}B \\ \downarrow J_{n-1}\rho_f^n & & \downarrow \rho_f^n & & \downarrow \\ J_{n-1}I_n[f] & \xrightarrow[\mu_{I_n[f]}^{n-1}]{} & I_n[f] & \twoheadrightarrow & I_{n-1}I_n[f] \end{array} \quad \begin{array}{c} \text{(iv)} \\ \text{(v)} \end{array}$$

We must prove that the right hand column is short exact. By the strongly $\mathcal{E}^{n-1}$ -Birkhoff property of I_{n-1} , the square (v) is an n -extension, hence $J_{n-1}\rho_f^n$ is an $(n-1)$ -extension. In particular, it is a regular epimorphism. Since, furthermore,

the square (iv) is a pullback (because $J_n[f] \subseteq J_{n-1}B$), the left hand column is short exact, hence, by applying the 3×3 -Lemma in the category $\text{Arr}^{n-1}\mathcal{A}$, so is the right hand one. $\square$

Lemma 6.3. *Let $\mathbb{G}$ be a comonad on $\mathcal{A}$ like in Subsection 5.2. For every $n \geq 1$, given an n -extension $f: B \rightarrow A$, there exists an isomorphism $K[I_n\mathbb{G}_nf] \cong K[I_{n-1}\mathbb{G}_nf]$ in $\mathcal{S}\text{Arr}^{n-1}\mathcal{A}$.*

Proof. $I_n\mathbb{G}_nf$, which consists degreeewise of central extensions with an $\mathcal{E}^n$ -projective codomain, is degreewise trivial by Proposition 4.5. This means that its kernel $K[I_n\mathbb{G}_nf]$ is isomorphic to $K[I_{n-1}I_n\mathbb{G}_nf]$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K[I_n\mathbb{G}_nf] & \xrightarrow{\quad} & I_n[\mathbb{G}_nf] & \xrightarrow{I_n\mathbb{G}_nf} & \mathbb{G}_{n-1}A \longrightarrow 0 \\ & & \cong \downarrow & & \eta_{I_n[\mathbb{G}_nf]}^{n-1} \downarrow & \lrcorner & \downarrow \eta_{\mathbb{G}_{n-1}A}^{n-1} \\ 0 & \longrightarrow & K[I_{n-1}I_n\mathbb{G}_nf] & \xrightarrow{\quad} & I_{n-1}I_n[\mathbb{G}_nf] & \xrightarrow{I_{n-1}I_n\mathbb{G}_nf} & I_{n-1}\mathbb{G}_{n-1}A \longrightarrow 0 \end{array}$$

By Lemma 6.2, the bottom short exact sequence is isomorphic to

$$0 \longrightarrow K[I_{n-1}\mathbb{G}_nf] \xrightarrow{\quad} I_{n-1}\mathbb{G}_{n-1}B \xrightarrow{I_{n-1}\mathbb{G}_nf} I_{n-1}\mathbb{G}_{n-1}A \longrightarrow 0, \quad (\mathbf{H})$$

which implies that $K[I_n\mathbb{G}_nf] \cong K[I_{n-1}\mathbb{G}_nf]$. $\square$

Theorem 6.4. *For any $n \geq 1$, $k \geq 2$ and any presentation $f: B \rightarrow A$ of an $(n-1)$ -extension A ,*

$$H_k(f, I_n)_{\mathbb{G}_n} \cong (H_{k+1}(A, I_{n-1})_{\mathbb{G}_{n-1}} \rightarrow 0).$$

Proof. We need to compute the homology of

$$I_n\mathbb{G}_nf: I_n[\mathbb{G}_nf] \rightarrow \mathbb{G}_{n-1}A,$$

a simplicial object in $\text{CExt}_{\mathcal{B}}^n\mathcal{A}$ —the image through I_n of the simplicial resolution $\mathbb{G}_nf: \mathbb{G}_{n-1}B \rightarrow \mathbb{G}_{n-1}A$ in $\text{Ext}^n\mathcal{A}$ of the extension f of A . By definition, the homology objects $H_{k-1}I_n\mathbb{G}_nf$ may be computed degreeewise, i.e.,

$$H_{k-1}I_n\mathbb{G}_nf \cong (H_{k-1}I_n[\mathbb{G}_nf] \rightarrow H_{k-1}\mathbb{G}_{n-1}A).$$

Note that $H_{i-1}\mathbb{G}_{n-1}A = 0$ for all $i \geq 2$; by the long exact homology sequence [18, Corollary 5.7], this implies that $H_{k-1}I_n[\mathbb{G}_nf]$ is isomorphic to $H_{k-1}K[I_n\mathbb{G}_nf]$. By Lemma 6.3, we know that the simplicial objects $K[I_n\mathbb{G}_nf]$ and $K[I_{n-1}\mathbb{G}_nf]$ are isomorphic. Using the long exact homology sequence associated with Sequence **H** and the fact that B is extension-projective (so that $\mathbb{G}_{n-1}B \rightarrow B$ is contractible), the desired isomorphism

$$H_{k-1}I_n[\mathbb{G}_nf] \cong H_kI_{n-1}\mathbb{G}_{n-1}A$$

is obtained. $\square$

7. THE HOPF FORMULA FOR THE SECOND HOMOLOGY OBJECT

In this section we give a direct proof of the Hopf formula from [18], which describes the second homology object in terms of “generalized commutators”.

Recall [18, Corollary 3.10] that $H_0S = \text{Coeq}[\partial_0, \partial_1: S_1 \rightarrow S_0]$ for every simplicial object S in a semi-abelian category $\mathcal{A}$ —a consequence of the fact that in $\mathcal{A}$, a regular epimorphism is a cokernel of its kernel.

Lemma 7.1. *Let $f: B \longrightarrow A$ be an n -extension; then $H_0 I_n \mathbb{G}_n f \cong I_n f$.*

Proof. The diagram

$$\begin{array}{ccccc} I_n[G^2 f] & \xrightleftharpoons[I_n[\epsilon_{Gf}]]{I_n[G\epsilon_f]} & I_n[Gf] & \xrightarrow{I_n[\epsilon_f]} & I_n[f] \\ I_n G^2 f \downarrow & & \downarrow I_n Gf & & \downarrow I_n f \\ G^2 A & \xrightleftharpoons[\epsilon_{GA}]{G\epsilon_A} & GA & \xrightarrow{\epsilon_A} & A \end{array}$$

is a coequalizer, because the square

$$\begin{array}{ccc} GB & \xrightarrow{\epsilon_B} & B \\ \rho_{Gf}^n \downarrow & & \downarrow \rho_f^n \\ I_n[Gf] & \xrightarrow{I_n[\epsilon_f]} & I_n[f] \end{array}$$

is a pushout: this follows from the strongly $\mathcal{E}^n$ -Birkhoff property of I_n . $\square$

Theorem 7.2. [18] *Let $f: B \longrightarrow A$ be a 1-presentation; then*

$$H_2(A, I)_{\mathbb{G}} \cong \frac{JB \cap K[f]}{J_1[f]}.$$

Proof. Note that $J_1[f]$ is a normal subobject of $JB \cap K[f]$ by definition. Since B is extension-projective, $I\mathbb{G}B$ is contractible, hence the short exact sequence of simplicial objects

$$0 \longrightarrow K[I\mathbb{G}f] \xrightarrow{\text{Ker } I\mathbb{G}f} I\mathbb{G}B \xrightarrow{I\mathbb{G}f} I\mathbb{G}A \longrightarrow 0$$

yields an exact homology sequence

$$0 \longrightarrow H_2(A, I)_{\mathbb{G}} = H_1 I\mathbb{G}A \longrightarrow H_0 K[I\mathbb{G}f] \longrightarrow IB \xrightarrow{If} IA \longrightarrow 0.$$

Taking into account Lemma 6.3, we thus get a short exact sequence

$$0 \longrightarrow H_2(A, I)_{\mathbb{G}} \longrightarrow H_0 K[I_1 \mathbb{G}f] \longrightarrow K[I f] \longrightarrow 0. \quad (\mathbf{I})$$

Now, using the long exact homology sequence induced by

$$0 \longrightarrow K[I_1 \mathbb{G}f] \longrightarrow I_1[\mathbb{G}f] \xrightarrow{I_1 \mathbb{G}f} \mathbb{G}A \longrightarrow 0$$

we find that

$$H_0 K[I_1 \mathbb{G}f] = K[H_0 I_1 \mathbb{G}f].$$

Hence, by the foregoing lemma,

$$H_0 K[I_1 \mathbb{G}f] = K[I_1 f].$$

By the 3×3 -Lemma,

$$K\left[I_1 f: \frac{B}{J_1[f]} \longrightarrow A\right] = \frac{K[f]}{J_1[f]}$$

and

$$K\left[If: \frac{B}{JB} \longrightarrow \frac{A}{JA}\right] = \frac{K[f]}{JB \cap K[f]}.$$

The theorem now follows from applying Noether's First Isomorphism Theorem (Theorem 4.3.10 in [3], a direct consequence of the 3×3 -Lemma) to **I**. $\square$

This also works for centralization of n -extensions:

Proposition 7.3. *Consider $n \in \mathbb{N}$ and $f: B \rightarrow A$, a presentation of an n -extension A . Then*


$$H_2(A, I_n)_{\mathbb{G}_n} \cong \frac{J_n B \cap K[f]}{J_{n+1}[f]} :$$

$H_2(A, I_n)_{\mathbb{G}_n}$ is the direct image of $K[f] \cap J_n B$ along $\rho_f^{n+1}: B \rightarrow I_{n+1}[f]$. $\square$

Remark 7.4. In particular, the expressions on the right hand side of these Hopf formulae are *Baer invariants* [19, 17]: they are independent of the chosen presentation of A .

Remark 7.5. Note that the objects in this Hopf formula are in $\text{Arr}^n \mathcal{A}$.

Notation 7.6. The functor category $\text{Arr}^n \mathcal{A} = \text{Hom}(2^n, \mathcal{A})$ may with advantage be described in the following way [11]. Let $\langle n \rangle$ denote the set $\{1, \dots, n\}$, $\langle 0 \rangle = \emptyset$. Then the category 2^n is isomorphic to the power-set $\mathcal{P}\langle n \rangle$, the set of subsets of $\langle n \rangle$, ordered by inclusion. This means that an inclusion $A \subseteq B$ in $\langle n \rangle$ corresponds to a map $\iota_B^A: A \rightarrow B$ in 2^n . We write f_A for $f(\iota_A^A)$ and f_B^A for $f(\iota_B^A)$. When, in particular, A is $\emptyset$ and B is a singleton $\{i\}$, we write $f_i = f_{\{i\}}^{\emptyset}$. For instance, $f_{\emptyset}$ is the “initial object” of the cube



and f_i , for $i \in \langle 3 \rangle$, is any of its three initial ribs.

Let us now write the Hopf formula using the notations just defined. By Notation 4.9, the kernel $K[\eta_B^n] = J_n B$ is written $\Psi^n L_n[B]$; it follows that the intersection $J_n B \cap K[f]$ is $\Psi^n(L_n[B] \cap K[f_{n+1}])$. Also, the kernel $J_{n+1}[f]$ is $\Psi^n L_{n+1}[f]$, and hence

$$H_2(A, I_n)_{\mathbb{G}_n} \cong \Psi^n \frac{L_n[B] \cap K[f_{n+1}]}{L_{n+1}[f]}.$$

Notation 7.7. An n -extension f naturally gives rise to k -extensions for all $0 \leq k \leq n$. We make the following choice of $\text{Ext}_k f \in \text{Ext}^k \mathcal{A}$: put $\text{Ext}_n f = f$ and $\text{Ext}_{k-1} f = \text{cod Ext}_k f$ ($0 < k \leq n$). It is easily seen that $\text{Ext}_k f$ may be obtained from f by precomposing with the functor $2^k \rightarrow 2^n$ determined by $A \mapsto A \cup (\langle n \rangle \setminus \langle k \rangle)$.

8. THE HIGHER HOPF FORMULAE

Let us start this section on the higher Hopf formulae by explaining the formula for the third homology object in the special case of canonically chosen presentations;

the others—where $n > 3$ or presentations are chosen non-canonically—work by essentially the same principle.

Suppose that $\mathcal{A}$ is a semi-abelian category and $\mathcal{B}$ is Birkhoff in $\mathcal{A}$. Let $\mathbb{G}$ be a comonad on $\mathcal{A}$ derived from an adjunction $\mathbf{F}$ which satisfies the conditions (1)–(2) from Subsection 5.2. Consider an object A in $\mathcal{A}$. By Theorem 6.4,

$$(H_3(A, I)_{\mathbb{G}} \longrightarrow 0) \cong H_2(\epsilon_A, I_1)_{\mathbb{G}_1},$$

i.e., in order to compute the third homology object of A with respect to I , it suffices to compute the second homology extension with respect to I_1 of the canonical presentation $\epsilon_A: GA \longrightarrow A$. This may be done using Proposition 7.3:

$$H_2(\epsilon_A, I_1)_{\mathbb{G}_1} \cong \rho_{\epsilon_A}^2(K[\epsilon_A^2] \cap K[\eta_{G\epsilon_A}^1]).$$

Here $\epsilon_A^2: G\epsilon_A \longrightarrow \epsilon_A$ is the canonical double extension of A , which has as kernel $K[\epsilon_A^2]$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & K[\epsilon_{GA}] & \twoheadrightarrow & G^2 A & \xrightarrow{\epsilon_{GA}} & GA \longrightarrow 0 \\ & & \downarrow K[\epsilon_A^2] & & \downarrow G\epsilon_A & & \downarrow \epsilon_A \\ 0 & \longrightarrow & K[\epsilon_A] & \twoheadrightarrow & GA & \xrightarrow{\epsilon_A} & A \longrightarrow 0. \end{array}$$

The other factor in the intersection is the kernel $K[\eta_{G\epsilon_A}^1]$ of the right hand side square in the next diagram.

$$\begin{array}{ccccccc} 0 & \longrightarrow & J_1[G\epsilon_A] & \twoheadrightarrow & G^2 A & \longrightarrow & I_1[G\epsilon_A] \longrightarrow 0 \\ & & \downarrow J_1 G\epsilon_A = K[\eta_{G\epsilon_A}^1] & & \downarrow G\epsilon_A & & \downarrow I_1 G\epsilon_A \\ 0 & \longrightarrow & 0 & \twoheadrightarrow & GA & \xlongequal{\quad} & GA \longrightarrow 0 \end{array}$$

The map $\rho_{\epsilon_A}^2: G\epsilon_A \longrightarrow I_2\epsilon_A^2$ may be pictured as

$$\begin{array}{ccc} G^2 A & \xrightarrow{\eta_{\epsilon_A^2}^2} & I_2[\epsilon_A^2]_0 \\ \downarrow G\epsilon_A & & \downarrow \\ GA & \xlongequal{\quad} & GA. \end{array}$$

Hence the direct image of $K[\eta_{G\epsilon_A}^1] \cap K[\epsilon_A^2]$ along $\rho_{\epsilon_A}^2$ is the front square of the cube

$$\begin{array}{ccccc} & & J_1[G\epsilon_A] \cap K[\epsilon_{GA}] & \twoheadrightarrow & G^2 A \\ & \swarrow & \downarrow & \swarrow & \downarrow G\epsilon_A \\ J_1[G\epsilon_A] \cap K[\epsilon_{GA}] & \xrightarrow{K[\rho_{\epsilon_A^2}^2]} & I_2[\epsilon_A^2]_0 & & \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \xrightarrow{\quad} & 0 & \longrightarrow & GA \\ \downarrow & & \downarrow & & \downarrow \\ 0 & \xrightarrow{\quad} & 0 & \longrightarrow & GA. \end{array}$$

If we write $L_2[\epsilon_A^2] = K[\rho_{\epsilon_A^2}^2]$, thus we get that

$$H_3(A, I)_{\mathbb{G}} \cong \frac{J_1[G\epsilon_A] \cap K[\epsilon_{GA}]}{L_2[\epsilon_A^2]}.$$

Using the Hopf formula for the second homology object, this may be rewritten as

$$H_3(A, I)_{\mathbb{G}} \cong \frac{JG^2A \cap K[G\epsilon_A] \cap K[\epsilon_{GA}]}{L_2[\epsilon_A^2]}, \quad (\mathbf{J})$$

because GA is projective, and hence $H_2(GA, I)_{\mathbb{G}} = 0$, which implies that

$$J_1[G\epsilon_A] \cong JG^2A \cap K[G\epsilon_A].$$

Modulo the more abstract denominator, this latter form **J** of the formula is how it occurs in Brown and Ellis's paper [11]. The main difference is that here, the formula is also valid for other categories $\mathcal{A}$ than the category $\mathbf{Gp}$ of groups, and other Birkhoff subcategories $\mathcal{B}$ than the category $\mathbf{Ab}$ of abelian groups.

Following the lines of the method due to Janelidze as set out in [26] and sketched above, we now show that also the higher Hopf formulae from [11] may be generalized to arbitrary semi-abelian categories.

We speak of an n -presentation f of an object A in $\mathcal{A}$ if f is an n -presentation and if $f_{\langle n \rangle} = A$ or, in other words, if $\text{Ext}_0 f = A$.

Theorem 8.1. *Let f be an n -presentation of an object A of $\mathcal{A}$. Then*

$$H_{n+1}(A, I)_{\mathbb{G}} \cong \frac{Jf_{\emptyset} \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{L_n[f]}. \quad (\mathbf{K})$$

*In particular, the expression on the right hand side of Formula **K** is a (higher) Baer invariant: independent of the chosen n -presentation of A .*

Proof. The formula holds by induction on n . Consider the following chain of isomorphisms (cf. Notation 7.7):

$$\begin{aligned} \Psi^{n-1} H_{n+1}(A, I)_{\mathbb{G}} &\cong \Psi^{n-2} H_n(\text{Ext}_1 f, I_1)_{\mathbb{G}_1} \\ &\cong \dots \cong \Psi H_3(\text{Ext}_{n-2} f, I_{n-2})_{\mathbb{G}_{n-2}} \\ &\cong H_2(\text{Ext}_{n-1} f, I_{n-1})_{\mathbb{G}_{n-1}} \\ &\cong \Psi^{n-1} \frac{L_{n-1}[\text{dom } f] \cap K[f_n]}{L_n[f]} \\ &\cong \Psi^{n-1} \frac{Jf_{\emptyset} \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{L_n[f]}. \end{aligned}$$

The first isomorphism follows from Theorem 6.4. Repeatedly applying this theorem gives the expression containing H_2 . Proposition 7.3 and Notation 7.6 account for the next isomorphism, and the last one follows by the induction hypothesis, the Hopf formula for $H_n(f_{\langle n \rangle \setminus \{n\}}, I)_{\mathbb{G}}$, because $\text{dom } f$ is an $(n-1)$ -presentation of $f_{\langle n \rangle \setminus \{n\}}$. Indeed, $f_{\langle n \rangle \setminus \{n\}}$ is a projective object, and hence $H_n(f_{\langle n \rangle \setminus \{n\}}, I)_{\mathbb{G}} = 0$, which implies that an isomorphism

$$L_{n-1}[\text{dom } f] \cong Jf_{\emptyset} \cap \bigcap_{i \in \langle n-1 \rangle} K[f_i]$$

exists. □

Remark 8.2. In particular, the formula in the Theorem above shows that the homology objects are independent of the chosen comonad on $\mathcal{A}$.

When a canonical presentation of A is chosen, one obtains the following formula.

Corollary 8.3. *For every $n \geq 1$, an isomorphism*

$$H_{n+1}(A, I)_{\mathbb{G}} \cong \frac{JG^n A \cap \bigcap_{i \in [n-1]} K[G^{n-1-i} \epsilon_{G^i A}]}{L_n[\epsilon_A^n]}$$

exists.

□

9. EXAMPLES

In this final section we consider some specific cases of Theorem 8.1 and give explicit descriptions of the right hand side of the Hopf formulae. In particular, we will explain why Theorem 8.1 gives Brown and Ellis's formulae [11] in the case where $\mathcal{A}$ is $\mathbf{Gp}$, the variety of groups and $\mathcal{B}$ is $\mathbf{Ab}$, the subvariety of all abelian groups. More generally, if $\mathcal{A}$ is $\mathbf{Gp}$ and $\mathcal{B}$ is $\mathbf{Nil}_k$, the subvariety of k -nilpotent groups (for some positive integer k), then we obtain the formulae due to Donadze, Inassaridze and Porter [15]. Furthermore, we find new formulae when $\mathcal{B} = \mathbf{Sol}_k$, the subvariety of k -solvable groups. Also, we explain that, by using the same arguments, similar formulae can be obtained from Theorem 8.1 when $\mathcal{A}$ is, e.g., the variety of rings or the variety of Lie algebras. Finally, we shall consider the case where $\mathcal{A}$ is $\mathbf{PXMod}$, the variety of precrossed modules, and $\mathcal{B}$ is $\mathbf{XMod}$, the subvariety of crossed modules.

From now on, we shall drop $\mathbb{G}$ in the notation for homology. This is justified by Remark 8.2 and by the fact that we shall be dealing with varieties and their canonical comonad.

9.1. Groups vs. abelian groups. Let us consider the Hopf formulae 8.1 in the particular case where $\mathcal{A} = \mathbf{Gp}$ is the variety of groups and $\mathcal{B} = \mathbf{Ab}$ the subvariety of abelian groups. In order to simplify the arguments below, we define the commutator of two (not necessarily normal) subgroups A and B of a group G as the *normal* subgroup of G generated by all elements $[a, b] = aba^{-1}b^{-1}$, with $a \in A$ and $b \in B$. Note that the reflector $I = \mathbf{ab}: \mathbf{Gp} \rightarrow \mathbf{Ab}$ is given by $\mathbf{ab}(G) = G/[G, G]$. It is known (see, e.g., [20] and Example 4.2) that in this situation $L_1[f] = [K[f], B]$ for any extension $f: B \rightarrow A$, hence when f is a presentation, Formula 8.1 yields, for $n = 1$, the classical Hopf formula

$$H_2(G, \mathbf{ab}) \cong \frac{[B, B] \cap K[f]}{[K[f], B]}.$$

We are now going to show that our formulae for $H_n(G, \mathbf{ab})$ coincide with those of Brown and Ellis, also when $n \geq 2$. In particular, we shall prove that, for all $n \geq 0$ and any n -extension f ,

$$L_n[f] = \prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right], \quad (\mathbf{L})$$

where it is understood that $\bigcap_{i \in \emptyset} K[f_i] = f_{\emptyset}$.

Let us recall the following *Witt-Hall identities*, valid in any group G , for any elements $a, a_1, a_2, b, b_1, b_2 \in G$:

$$[a, b_1 b_2] = [a, b_1] b_1 [a, b_2] b_1^{-1} \quad (\mathbf{M})$$

$$[a_1 a_2, b] = a_1 [a_2, b] a_1^{-1} [a_1, b]. \quad (\mathbf{N})$$

Let us write $A \cdot B$ for the product of subgroups A and B of a group G . The identities **M** and **N** imply, for all groups G and subgroups A, A_1, A_2, B, B_1 and B_2 of G that

$$[A, B_1 \cdot B_2] = [A, B_1] \cdot [A, B_2]$$

and

$$[A_1 \cdot A_2, B] = [A_1, B] \cdot [A_2, B].$$

For a group G , we denote by $\Delta_G = (1_G, 1_G): G \longrightarrow G \times G$ the *diagonal* of G : the smallest internal equivalence relation on G .

Then, in particular, we get

Lemma 9.1. *Suppose G is a group, M and N are normal subgroups of G and B is a subgroup of $G \times G$. Then the following identity holds:*

$$[M \times_{\frac{M}{M \cap N}} M, B] = [0 \times (M \cap N), B] \cdot [\Delta_M, B].$$

Proof. This follows from the above, by taking into account that

$$M \times_{\frac{M}{M \cap N}} M = 0 \times (M \cap N) \cdot \Delta_M,$$

for any normal subgroups M and N of a group G . $\square$

Let us then prove the identity **L**, for all $n \geq 0$ and any n -extension f . We do this by induction. For $n = 0$, we have

$$L_0[A] = JA = [A, A] = \prod_{I \subseteq \emptyset} [A, A],$$

when $A = f$ is a 0-extension in $\mathbf{Gp}$. Now, assume that **L** holds for some $n - 1$. To show that **L** holds for n , it suffices to prove that

$$L_n[f] = 0 \Leftrightarrow \prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right] = 0, \quad (\mathbf{O})$$

for every n -extension f . Indeed, suppose the equivalence **O** holds. Recall that, for any group G and any subgroups A and B of G , $[A, B]$ is the smallest normal subgroup N of G such that

$$\left[\frac{A \cdot N}{N}, \frac{B \cdot N}{N} \right] = 0.$$

From this one deduces that

$$\prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right]$$

is the smallest normal subgroup N of $f_\emptyset$ satisfying

$$\prod_{I \subseteq \langle n \rangle} \left[\frac{\bigcap_{i \in I} K[f_i] \cdot N}{N}, \frac{\bigcap_{i \notin I} K[f_i] \cdot N}{N} \right] = 0.$$

Since $L_n[I_n f] = 0$, applying $\mathbf{O}$ to the n -extension $I_n f$ gives that

$$\prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} \frac{K[f_i]}{L_n[f]}, \bigcap_{i \notin I} \frac{K[f_i]}{L_n[f]} \right] = \prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[(I_n f)_i], \bigcap_{i \notin I} K[(I_n f)_i] \right] = 0,$$

hence

$$L_n[f] \supseteq \prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right].$$

The other inclusion follows similarly, now taking into account that I_n is a reflector.

Let us then prove the equivalence $\mathbf{O}$. Suppose $f: B \rightarrow A$ is an n -extension. It follows readily from the definition of $J_n = \Psi L_n$ that $L_n[f] = 0$ if and only if

$$\pi_1 L_{n-1}[R[f]] = \pi_2 L_{n-1}[R[f]],$$

where (π_1, π_2) denotes the kernel pair of $f_n: f_\emptyset \rightarrow f_{\{n\}}$. By the induction hypothesis,

$$L_{n-1}[R[f]] = \prod_{I \subseteq \langle n-1 \rangle} \left[\bigcap_{i \in I} K[R[f]_i], \bigcap_{i \notin I} K[R[f]_i] \right].$$

Recall from Notation 7.6 that $R[f]_A = R[f_{A \cup \{n\}}^A]$ for all $A \subseteq \langle n-1 \rangle$. In particular, $R[f]_\emptyset = R[f_n]$ and $R[f]_{\{i\}} = R[f_{\{i,n\}}^{\{i\}}]$. Furthermore, $R[f]_i$ is such that the following diagram commutes.

$$\begin{array}{ccccc} R[f]_\emptyset & \rightrightarrows & f_\emptyset & \xrightarrow{f_n} & f_{\{n\}} \\ R[f]_i \downarrow & & f_i \downarrow & & \downarrow f_{\{i,n\}}^{\{i\}} \\ R[f]_{\{i\}} & \rightrightarrows & f_{\{i\}} & \xrightarrow{f_{\{i,n\}}^{\{i\}}} & f_{\{i,n\}} \end{array}$$

Since the right hand square is a regular pushout, Proposition 3.3 implies that

$$K[f_{\{i,n\}}^{\{i\}}] = \frac{K[f_i]}{K[f_n] \cap K[f_i]}.$$

Consequently,

$$K[R[f]_i] = K[f_i] \times_{\frac{K[f_i]}{K[f_n] \cap K[f_i]}} K[f_i].$$

It follows that

$$L_{n-1}[R[f]] = \prod_{I \subseteq \langle n-1 \rangle} \left[\bigcap_{i \in I} K[f_i] \times_{\frac{K[f_i]}{K[f_i] \cap K[f_n]}} K[f_i], \bigcap_{i \notin I} K[f_i] \times_{\frac{K[f_i]}{K[f_i] \cap K[f_n]}} K[f_i] \right].$$

Moreover note that, for all $I \subseteq \langle n \rangle$, we have

$$\bigcap_{i \in I} K[f_i] \times_{\frac{K[f_i]}{K[f_i] \cap K[f_n]}} K[f_i] = \bigcap_{i \in I} K[f_i] \times_{\frac{\bigcap_{i \in I} K[f_i]}{\bigcap_{i \in I} K[f_i] \cap K[f_n]}} \bigcap_{i \in I} K[f_i].$$

Now, applying Lemma 9.1, we find that

$$\begin{aligned}
L_{n-1}[R[f]] &= \prod_{I \subseteq \langle n-1 \rangle} \left[0 \times \bigcap_{i \in I} K[f_i] \cap K[f_n], 0 \times \bigcap_{i \notin I} K[f_i] \cap K[f_n] \right] \\
&\quad \cdot \left[0 \times \bigcap_{i \in I} K[f_i] \cap K[f_n], \Delta_{\bigcap_{i \notin I} K[f_i]} \right] \\
&\quad \cdot \left[\Delta_{\bigcap_{i \in I} K[f_i]}, 0 \times \bigcap_{i \notin I} K[f_i] \cap K[f_n] \right] \\
&\quad \cdot \left[\Delta_{\bigcap_{i \in I} K[f_i]}, \Delta_{\bigcap_{i \notin I} K[f_i]} \right].
\end{aligned}$$

Consequently, $\pi_1 L_{n-1}[R[f]] = \pi_2 L_{n-1}[R[f]]$ if and only if

$$\left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \cap K[f_n] \right] = \{1\},$$

for all $I \subseteq \langle n-1 \rangle$, i.e., if and only if

$$\prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right] = \{1\},$$

which is exactly what we wanted. We conclude that the equality **L** holds for all $n \geq 0$. Thus Theorem 8.1 induces Brown and Ellis's formulae:

Theorem 9.2. [11] *For any $n \geq 1$ and any n -presentation f of a group G , an isomorphism*

$$H_{n+1}(G, \text{ab}) \cong \frac{[f_\emptyset, f_\emptyset] \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{\prod_{I \subseteq \langle n \rangle} \left[\bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right]}$$

exists. □

9.2. Groups vs. k -nilpotent groups. Recall that a group A is k -nilpotent (= nilpotent of class at most k) if and only if $Z_k A = \{1\}$, where $Z_k A$ is the k -th term in the *descending central series* of A defined by

$$\begin{aligned}
Z_1 A &= [A, A], \\
Z_2 A &= [[A, A], A], \\
\ldots &= \ldots \\
Z_k A &= [Z_{k-1} A, A].
\end{aligned}$$

Let us now consider the Hopf formulae 8.1 in the situation where $\mathcal{A} = \mathbf{Gp}$ and $\mathcal{B} = \mathbf{Nil}_k$, the subvariety of $\mathbf{Gp}$ of all k -nilpotent groups (for some positive integer k). Let us write nil_k for the reflector $\mathbf{Gp} \longrightarrow \mathbf{Nil}_k$, which sends a group A to the quotient $A/Z_k A$.

The arguments used above in Subsection 9.1 can easily be adapted to show that Lemma 9.1 implies, in this situation, that

$$L_n[f] = \prod_{I_1 \cup \ldots \cup I_k = \langle n \rangle} \left[\ldots \left[\left[\bigcap_{i \in I_1} K[f_i], \bigcap_{i \in I_2} K[f_i] \right], \bigcap_{i \in I_3} K[f_i] \right], \ldots \right],$$

for any $n \geq 0$ and any n -extension of groups f . Consequently, the Hopf formulae 8.1 yield

Theorem 9.3. [15] *For any $n \geq 1$ and any n -presentation f of a group G , an isomorphism*

$$H_{n+1}(G, \text{nil}_k) \cong \frac{[\dots [[f_\emptyset, f_\emptyset], f_\emptyset], \dots] \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{\prod_{I_1 \cup \dots \cup I_k = \langle n \rangle} \left[\dots \left[\left[\bigcap_{i \in I_1} K[f_i], \bigcap_{i \in I_2} K[f_i] \right], \bigcap_{i \in I_3} K[f_i] \right], \dots \right]}$$

exists. $\square$

9.3. Groups vs. k -solvable groups. Recall that a group A is k -solvable (= solvable of class at most k) if and only if $D_k A = \{1\}$, where $D_k A$ is the k -th term in the *derived series* of A defined by

$$\begin{aligned} D_1 A &= [A, A], \\ D_2 A &= [[A, A], [A, A]], \\ \dots &= \dots \\ D_k A &= [D_{k-1} A, D_{k-1} A]. \end{aligned}$$

Let us consider the Hopf formulae 8.1 in the case where $\mathcal{A} = \text{Sol}_k$, the subvariety of $\mathbf{Gp}$ of all k -solvable groups (for some positive integer k). We write sol_k for the reflector $\mathbf{Gp} \rightarrow \text{Sol}_k$, which sends a group A to the quotient $A/D_k A$. In order to express a formula for $L_n[\cdot]$, it is useful to introduce also the following notation:

Given normal subgroups $X_1, X_2, X_3, \dots$ of a group A , one defines inductively

$$\begin{aligned} D_1(X_1, X_2) &= [X_1, X_2], \\ D_2(X_1, X_2, X_3, X_4) &= [[X_1, X_2], [X_3, X_4]], \\ \dots &= \dots \\ D_k(X_1, \dots, X_{2^k}) &= [D_{k-1}(X_1, \dots, X_{2^{k-1}}), D_{k-1}(X_{2^{k-1}+1}, \dots, X_{2^k})]. \end{aligned}$$

Thus, in this situation, $J(A) = D_k(A, \dots, A)$. Furthermore, the arguments of Subsection 9.1 are easily adapted to show that Lemma 9.1 implies

$$L_n[f] = \prod_{I_1 \cup \dots \cup I_{2^k} = \langle n \rangle} D_k \left(\bigcap_{i \in I_1} K[f_i], \dots, \bigcap_{i \in I_{2^k}} K[f_i] \right),$$

for any $n \geq 0$ and any n -extension of groups f . Consequently, Theorem 8.1 yields the following formulae:

Theorem 9.4. *For any $n \geq 1$ and any n -presentation f of a group G , an isomorphism*

$$H_{n+1}(G, \text{sol}_k) \cong \frac{D_k(f_\emptyset, \dots, f_\emptyset) \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{\prod_{I_1 \cup \dots \cup I_{2^k} = \langle n \rangle} D_k(\bigcap_{i \in I_1} K[f_i], \dots, \bigcap_{i \in I_{2^k}} K[f_i])}$$

exists. $\square$

9.4. Some additional examples. The reader will find it easy to further adapt the arguments used above to prove that Theorem 8.1 yields similar formulae, for example in the variety of non-unital rings (where the role of commutator is played by the product of subrings) or in the variety of Lie algebras (the Lie bracket plays the role of commutator). The example of precrossed modules versus crossed modules, however, deserves special attention.

9.5. Precrossed modules vs. crossed modules. Recall that a *precrossed module* (C, G, ∂) is a group homomorphism $\partial: C \longrightarrow G$ equipped with a (left) group action of G on C , such that

$$\partial(gc) = g\partial(c)g^{-1}$$

for all $g \in G$ and $c \in C$. A morphism $f: (C, G, \partial) \longrightarrow (D, H, \epsilon)$ of precrossed modules is a pair of group homomorphisms $f_1: C \longrightarrow D$ and $f_0: G \longrightarrow H$ which preserve the action and are such that $\epsilon \circ f_1 = f_0 \circ \partial$. Let us write $\mathbf{PXMod}$ for the category of precrossed modules and $\mathbf{XMod}$ for the category of *crossed modules*, where this latter is the full subcategory of $\mathbf{PXMod}$ whose objects (C, G, ∂) satisfy the *Peiffer condition*

$$\partial(c)c' = cc'c^{-1}$$

for all $c, c' \in C$. We will sometimes abbreviate the notation (C, G, ∂) to C .

It is well known that the category of precrossed modules is equivalent to a variety of Ω -groups (see, e.g., [31] [32] or [29]). Via this equivalence, $\mathbf{XMod}$ correspond to a subvariety of $\mathbf{PXMod}$. Hence, we can consider the Hopf formulae 8.1 in the case where $\mathcal{A} = \mathbf{PXMod}$ and $\mathcal{B} = \mathbf{XMod}$.

Recall that a *precrossed submodule* of a precrossed module (C, G, ∂) is a precrossed module (M, S, μ) such that M and S are, respectively, subgroups of C and G , and such that the action of S on M is a restriction of the action of G on C and μ a restriction of ∂ (in this case, we will write ∂ instead of μ). (M, S, ∂) is a *normal* precrossed submodule of (C, G, ∂) if, furthermore, M and S are, respectively, normal subgroups of C and G , and, for all $c \in C$, $g \in G$, $m \in M$, $s \in S$, one has ${}^g m \in M$ and ${}^s cc^{-1} \in M$. This is exactly the case when (M, S, ∂) is the kernel of some morphism $(C, G, \partial) \longrightarrow (D, H, \epsilon)$. Note that the quotient $(C, G, \partial) \longrightarrow (C, G, \partial)/(K, S, \partial)$ is a degreewise quotient

$$(q_M, q_S): (C, G, \partial) \longrightarrow (C/M, G/S, \bar{\partial}).$$

Here we have written q_M and q_S for the quotient homomorphisms $C \longrightarrow C/M$ and $G \longrightarrow G/S$, respectively. Note also that limits in $\mathbf{PXMod}$ are degreewise limits in $\mathbf{Gp}$.

Let (C, G, ∂) be a precrossed module. The *Peiffer commutator* of two precrossed submodules (M, S, ∂) and (N, T, ∂) of (C, G, ∂) is the normal subgroup of the group $M \cdot N$, generated by the *Peiffer elements* $\langle m, n \rangle = mn m^{-1} (\partial^m n)^{-1}$ and $\langle n, m \rangle = nm n^{-1} (\partial^n m)^{-1}$, with $m \in M$ and $n \in N$. We will denote it by $\langle (M, S, \partial), (N, T, \partial) \rangle$ or simply by $\langle M, N \rangle$. Note that, for all $m \in M$ and $n \in N$, we have that

$$\begin{aligned} \partial \langle m, n \rangle &= \partial m \partial n (\partial m)^{-1} (\partial^{\partial m} n)^{-1} \\ &= \partial m \partial n (\partial m)^{-1} (\partial m \partial n (\partial m)^{-1})^{-1} \\ &= 1. \end{aligned}$$

Hence we may consider $\langle M, N \rangle$ as the precrossed submodule $(\langle M, N \rangle, 0, \partial)$ of C .

The following simple identities, similar to **M** and **N**, are valid in any precrossed module (C, G, ∂) , for any elements $a, a_1, a_2, b, b_1, b_2 \in C$.

$$\langle a, b_1 b_2 \rangle = \langle a, b_1 \rangle^{\partial a} b_1 \langle a, b_2 \rangle (\partial^a b_1)^{-1} \quad (\mathbf{P})$$

$$\langle a_1 a_2, b \rangle = a_1 \langle a_2, b \rangle a_1^{-1} \langle a_1, \partial^{a_2} b \rangle \quad (\mathbf{Q})$$

For a precrossed module C , let us denote by $\Delta_C = (1_C, 1_C): C \longrightarrow C \times C$ the *diagonal* of C : the smallest internal equivalence relation on C .

Lemma 9.5. *Suppose (C, G, ∂) is a precrossed module, (M, S, ∂) and (N, T, ∂) are normal precrossed submodules of C , and $(B, P, \partial \times \partial)$ is a precrossed submodule of $C \times C$. When B is either a normal precrossed submodule of $C \times C$ or $B = \Delta_K$, for some normal precrossed submodule (K, R, ∂) of C , then the following identity holds:*

$$\langle M \times_{\frac{M}{M \cap N}} M, B \rangle = \langle 0 \times (M \cap N), B \rangle \cdot \langle \Delta_M, B \rangle.$$

Proof. Note that in both cases only one inclusion is not entirely trivial. We prove the other.

Let us first assume that B is a normal precrossed submodule of $C \times C$. Suppose that $m \in M$, $d \in M \cap N$ and $(b_1, b_2) \in B$, then by identity **Q**,

$$\langle (m, dm), (b_1, b_2) \rangle = \langle (1, d)(m, m), (b_1, b_2) \rangle \in \langle 0 \times (M \cap N), B \rangle \cdot \langle \Delta_M, B \rangle.$$

Similarly, by identity **P**, $\langle (b_1, b_2), (m, dm) \rangle \in \langle 0 \times (M \cap N), B \rangle \cdot \langle \Delta_M, B \rangle$.

Let us now assume that $B = \Delta_K$, for some normal precrossed submodule (K, R, ∂) of C . Take $m \in M$, $d \in M \cap N$ and $k \in K$. It follows from **P** that

$$\langle (k, k), (m, dm) \rangle = \langle (k, k), (1, d)(m, m) \rangle \in \langle 0 \times (M \cap N), B \rangle \cdot \langle \Delta_M, B \rangle.$$

Furthermore,

$$\begin{aligned} \langle (m, dm), (k, k) \rangle &= (\langle m, k \rangle, \langle dm, k \rangle) \\ &= (\langle m, k \rangle, d\langle m, k \rangle d^{-1} \langle d, \partial^m k \rangle) \\ &= (\langle m, k \rangle, d\langle m, k \rangle d^{-1} (1, \langle d, \partial^m k \rangle)). \end{aligned}$$

On the one hand,

$$(\langle m, k \rangle, d\langle m, k \rangle d^{-1}) = (1, d)(\langle m, m \rangle, (k, k))(1, d)^{-1} \in \langle \Delta_M, B \rangle.$$

On the other hand,

$$(1, \langle d, \partial^m k \rangle) = \langle (1, d), (\partial^m k, \partial^m k) \rangle \in \langle 0 \times (M \cap N), B \rangle. \quad \square$$

Using Lemma 9.5, the reader will find it easy to adapt the arguments of Section 9.1 to show that

$$L_n[f] = \left(\prod_{I \subseteq \langle n \rangle} \left\langle \bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right\rangle, 0, \partial \right),$$

for any $n \geq 0$ and any n -extension of precrossed modules f . Consequently, Theorem 8.1 becomes

Theorem 9.6. *For any $n \geq 1$ and any n -presentation f of a precrossed module (C, G, ∂) , an isomorphism*

$$H_{n+1}((C, G, \partial), \text{xmod}) \cong \frac{\langle f_\emptyset, f_\emptyset \rangle \cap \bigcap_{i \in \langle n \rangle} K[f_i]}{\prod_{I \subseteq \langle n \rangle} \left\langle \bigcap_{i \in I} K[f_i], \bigcap_{i \notin I} K[f_i] \right\rangle}$$

exists. □

INDEX OF NOTATIONS

$\text{Arr}^n \mathcal{A}$, 7	f_A , 24	$L_n[\cdot]$, 17
$\text{CExt}_{\mathcal{B}} \mathcal{A}$, 14	$\mathbb{G}$, 18	μ^n , 14
$\text{CExt}_{\mathcal{B}}^n \mathcal{A}$, 14	Γ , 5	$\text{NExt}_{\mathcal{B}}^n \mathcal{A}$, 14
$\mathcal{E}^n$, 9, 14	Γ_n , 14	Ψ^n , 12
ϵ , 18	H_n , 18	$\text{Reg}^n \mathcal{A}$, 8
ϵ^n , 21	$H_n(\cdot, I)_{\mathbb{G}}$, 18	ρ^n , 14
η , 17	I_n , 13	$\subseteq_n$, 14
η^n , 14	$I_n[\cdot]$, 14	$\mathcal{Z}^n$, 14
$\text{Ext}^n \mathcal{A}$, 9	J_n , 13	
$\text{Ext}_n f$, 24	$J_n[\cdot]$, 14	

REFERENCES

- [1] M. Barr, *Exact categories*, Exact categories and categories of sheaves, Lecture notes in mathematics, vol. 236, Springer, 1971, pp. 1–120.
- [2] M. Barr and J. Beck, *Homology and standard constructions*, Seminar on triples and categorical homology theory, Lecture notes in mathematics, vol. 80, Springer, 1969, pp. 245–335.
- [3] F. Borceux and D. Bourn, *Mal'cev, protomodular, homological and semi-abelian categories*, Mathematics and its Applications, vol. 566, Kluwer Academic Publishers, 2004.
- [4] F. Borceux and G. Janelidze, *Galois theories*, Cambridge Studies in Advanced Mathematics, vol. 72, Cambridge University Press, 2001.
- [5] D. Bourn, *Normalization equivalence, kernel equivalence and affine categories*, in Carboni et al. [13], pp. 43–62.
- [6] ———, *Mal'cev categories and fibration of pointed objects*, Appl. Categ. Struct. **4** (1996), 307–327.
- [7] ———, *3×3 lemma and protomodularity*, J. Algebra **236** (2001), 778–795.
- [8] ———, *The denormalized 3×3 lemma*, J. Pure Appl. Algebra **177** (2003), 113–129.
- [9] D. Bourn and G. Janelidze, *Protomodularity, descent, and semidirect products*, Theory Appl. Categ. **4** (1998), no. 2, 37–46.
- [10] ———, *Characterization of protomodular varieties of universal algebras*, Theory Appl. Categ. **11** (2003), no. 6, 143–147.
- [11] R. Brown and G. J. Ellis, *Hopf formulae for the higher homology of a group*, Bull. London Math. Soc. **20** (1988), 124–128.
- [12] A. Carboni, G. M. Kelly, and M. C. Pedicchio, *Some remarks on Maltsev and Goursat categories*, Appl. Categ. Struct. **1** (1993), 385–421.
- [13] A. Carboni, M. C. Pedicchio, and G. Rosolini (eds.), *Category Theory, Proceedings Como 1990*, Lecture notes in mathematics, vol. 1488, Springer, 1991.
- [14] B. Conrad, *Crossed n -fold extensions of groups, n -fold extensions of modules, and higher multipliers*, J. Pure Appl. Algebra **36** (1985), 225–235.
- [15] G. Donadze, N. Inassaridze, and T. Porter, *n -Fold Čech derived functors and generalised Hopf type formulas*, K-Theory **35** (2005), no. 3–4, 341–373.
- [16] T. Everaert, *An approach to non-abelian homology based on Categorical Galois Theory*, Ph.D. thesis, Vrije Universiteit Brussel, 2007.
- [17] T. Everaert and T. Van der Linden, *Baer invariants in semi-abelian categories I: General theory*, Theory Appl. Categ. **12** (2004), no. 1, 1–33.
- [18] ———, *Baer invariants in semi-abelian categories II: Homology*, Theory Appl. Categ. **12** (2004), no. 4, 195–224.
- [19] A. Fröhlich, *Baer-invariants of algebras*, Trans. Amer. Math. Soc. **109** (1963), 221–244.
- [20] J. Furtado-Coelho, *Homology and generalized Baer invariants*, J. Algebra **40** (1976), 596–609.
- [21] M. Gran and J. Rosický, *Semi-abelian monadic categories*, Theory Appl. Categ. **13** (2004), no. 6, 106–113.

- [22] M. Gran and V. Rossi, *Galois theory and double central extensions*, Homology, Homotopy and Appl. **6** (2004), no. 1, 283–298.
- [23] H. Hopf, *Fundamentalgruppe und zweite Bettische Gruppe*, Comment. Math. Helv. **14** (1942), 257–309.
- [24] G. Janelidze, *Pure Galois theory in categories*, J. Algebra **132** (1990), 270–286.
- [25] ———, *Precategories and Galois theory*, in Carboni et al. [13], pp. 157–173.
- [26] ———, *What is a double central extension? (The question was asked by Ronald Brown)*, Cah. Top. Géom. Diff. Catég. **XXXII** (1991), no. 3, 191–201.
- [27] ———, *Higher dimensional central extensions: A categorical approach to homology theory of groups*, Lecture at the International Category Theory Meeting CT95, Halifax, 1995.
- [28] G. Janelidze and G. M. Kelly, *Galois theory and a general notion of central extension*, J. Pure Appl. Algebra **97** (1994), 135–161.
- [29] G. Janelidze, L. Márki, and W. Tholen, *Semi-abelian categories*, J. Pure Appl. Algebra **168** (2002), 367–386.
- [30] G. Janelidze, M. Sobral, and W. Tholen, *Beyond Barr exactness: Effective descent morphisms*, Categorical Foundations: Special Topics in Order, Topology, Algebra and Sheaf Theory (M. C. Pedicchio and W. Tholen, eds.), Encyclopedia of Mathematics and its Applications, vol. 97, Cambridge University Press, 2004, pp. 359–405.
- [31] R. Lavendhomme and J. R. Roisin, *Cohomologie non abélienne de structures algébriques*, J. Algebra **67** (1980), 385–414.
- [32] J.-L. Loday, *Spaces with finitely many non-trivial homotopy groups*, J. Pure Appl. Algebra **24** (1982), 179–202.
- [33] A. S.-T. Lue, *Baer-invariants and extensions relative to a variety*, Proc. Camb. Phil. Soc. **63** (1967), 569–578.
- [34] D. G. Quillen, *Homotopical algebra*, Lecture notes in mathematics, vol. 43, Springer, 1967.
- [35] A. Rodicio, *Presentaciones libres y $H_{2n}(G)$* , Publ. Mat. Univ. Aut. Barcelona **30** (1986), 77–82.
- [36] R. Stöhr, *A generalized Hopf formula for higher homology groups*, Comment. Math. Helvetici **64** (1989), 187–199.

VAKGROEP WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM
E-mail address: `teveraer@vub.ac.be`

LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES, UNIVERSITÉ DU LITTORAL CÔTE
 D'OPALE, 50, RUE F. BUISSON, 62228 CALAIS, FRANCE
E-mail address: `gran@lmpa.univ-littoral.fr`

VAKGROEP WISKUNDE, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM
E-mail address: `tvdlinde@vub.ac.be`